

of inequalities in integer
45-162.

ight of a graph. RUTCOR
sity, New Brunswick, NJ

Updating the Hamiltonian Problem— A Survey

Ronald J. Gould

DEPARTMENT OF MATHEMATICS
AND COMPUTER SCIENCE
EMORY UNIVERSITY
ATLANTA, GEORGIA

ABSTRACT

This is intended as a survey article covering recent developments in the area of hamiltonian graphs, that is, graphs containing a spanning cycle. This article also contains some material on related topics such as traceable, hamiltonian-connected and pancyclic graphs and digraphs, as well as an extensive bibliography of papers in the area.

0. INTRODUCTION

The hamiltonian problem; determining when a graph contains a spanning cycle has long been fundamental in graph theory. Named for Sir William Rowan Hamilton, this problem traces its origins to the 1850s. Today, however, the flood of papers dealing with this subject and its many related problems is at its greatest; supplying us with new results as well as many new problems involving cycles and paths in graphs.

To many, including myself, any path or cycle question is really a part of this general area. Although it is difficult to separate many of these ideas, for the purpose of this article, I will concentrate my efforts on results on problems dealing with spanning cycles (the classic hamiltonian problem) on ordinary graphs. I shall not attempt to survey digraphs, the traveling salesman problem (see instead [107]), or any of its related questions. However, I shall mention a few related results. I shall further restrict my attention primarily to work done since the late 1970s; however, for completeness, I shall include some earlier work in several places. For an excellent general introduction to the hamiltonian problem, the reader should see the article by J. C. Bermond [23]. Those not familiar with this topic or with graphs in general are advised to begin there. Further background and related material

Journal of Graph Theory, Vol. 15, No. 2, 121-157 (1991)
© 1991 John Wiley & Sons, Inc. CCC 0364-9024/91/020121-37\$04.00

can be found in the following related survey articles: [31], [24], [108], [163], [51], [19], and [124].

This article concludes with a rather extensive list of references. I have also tried to include the *Math Reviews* reference whenever possible. I hope this will be of use to those interested in research problems in this field.

Throughout this article we will consider finite graphs $G = (V, E)$. We reserve n to denote the order ($|V|$) of the graph under consideration and q the size ($|E|$). A graph will be called *hamiltonian* if it contains a spanning cycle. Such a cycle will be called a *hamiltonian cycle*. If a graph G contains a spanning path it is termed a *traceable* graph and if G contains a spanning path joining any two of its vertices, then G is *hamiltonian-connected*. If G contains a cycle of each possible length l , $3 \leq l \leq n$, then G is said to be *pancyclic*. These are clearly closely linked ideas and by no means does this list exhaust the related concepts.

There are four fundamental results that I feel deserve special attention here—both for their contribution to the overall theory and for their affect on the development of the area. In many ways, these four results are the foundation of much of today's work.

Beginning with Dirac's theorem [53] in 1952, the approach taken to developing sufficient conditions for a graph to be hamiltonian usually involved some sort of edge density condition; providing enough edges to overcome any obstructions to the existence of a hamiltonian cycle. Dirac saw a natural method for supplying the necessary edges, using the minimum degree $\delta(G)$.

Theorem 0.1 [53]. If G is a graph of order n such that $\delta(G) \geq n/2$, then G is hamiltonian.

Dirac's theorem was followed by that of Ore [132]. Ore's theorem relaxed Dirac's condition and extended the methods for controlling the degrees of the vertices in the graph.

Theorem 0.2 [132]. If G is a graph of order n such that $\deg x + \deg y \geq n$, for every pair of nonadjacent vertices $x, y \in V$, then G is hamiltonian.

This relaxation stimulated a string of subsequent refinements (see [44] or [23] for more details), culminating in the classic work of Bondy and Chvátal [33] concerning stability and closure. In [33], as in Ore's [132] motivating work, independent (mutually nonadjacent) vertices whose degree sum is at least n are fundamental. The following notation will be useful:

$$\sigma_k(G) = \min \left\{ \sum_{i=1}^k \deg v_i \mid \{v_1, v_2, \dots, v_k\} \text{ is independent in } G \ (k \geq 2) \right\}.$$

In [33], Bondy and Chvátal extended Ore's theorem in a very useful way. Define the *k-degree closure* of G , denoted $C_k(G)$, as the graph obtained

by recursively joining pairs of nonadjacent vertices whose degree sum is at least k , until no such pair remains. The following:

Theorem 0.3 [33]. A graph G is hamiltonian if and only if $C_k(G)$ is hamiltonian.

Theorem 0.3 provides an interesting generalization. We no longer need to verify that each vertex has degree at least $n/2$, but rather, only a certain edge density is recognizable as being hamiltonian. If a graph is hamiltonian, your chances should improve as you add edges. In forming the degree closure, examples for all possible values of k are given. No help in deciding if the original graph is hamiltonian may be the complete graph.

This idea led naturally to the definition of *k-degree closure* for all graphs. A graph is *k-degree stable* if, for all graphs H with the same property \hat{P} and $\deg u + \deg v \geq k$, the degree closure of H is hamiltonian. Results established in [33] were the

- (i) The property of being hamiltonian.
- (ii) The property of being traceable.
- (iii) The property of containing a cycle of length l .

For other work related to the degree closure, see [170].

The fourth fundamental result concerns the independence number of a graph. Note the independence number of a graph is the maximum size of an independent set of vertices in G .

Theorem 0.4 [47]. If G is a graph of order n such that $\sigma_k(G) \geq n - k + 1$, then G is hamiltonian.

In the following sections, we will discuss the degree closure and inspired many others.

1. GENERALIZATIONS OF THEOREM 0.3

Many generalizations of Theorem 0.3 have been found. Bondy and Chvátal's [33] sharpened Dirac's condition. The degree closure property of the graph into the degree closure.

Theorem 1.1 [86]. If G is a 2-connected graph of order n and minimum degree $\delta(G) \geq \frac{1}{3}(n-1)$, then G is hamiltonian.

by recursively joining pairs of nonadjacent vertices whose degree sum is at least k , until no such pair remains. Their fundamental hamiltonian result is the following:

Theorem 0.3 [33]. A graph G of order n is hamiltonian if, and only if, $C_n(G)$ is hamiltonian.

Theorem 0.3 provides an interesting relaxation of Ore's condition. Now we no longer need to verify that each pair of nonadjacent vertices has degree sum at least n , but rather, only enough pairs to ensure that the closure is recognizable as being hamiltonian. Since the closure is hopefully a denser graph, your chances should improve. However, the number of edges actually added in forming the degree closure can vary widely. It is easy to construct examples for all possible values from 0 to $\binom{n}{2} - q$. Thus, we might receive no help in deciding if the original graph is hamiltonian, or the degree closure may be the complete graph.

This idea led naturally to the following definition. Let \hat{P} be a property defined for all graphs of order n and let k be an integer. Then \hat{P} is said to be k -degree stable if, for all graphs G of order n , whenever $G + uv$ has property \hat{P} and $\deg u + \deg v \geq k$, then G has property \hat{P} . Among the results established in [33] were the following:

- (i) The property of being hamiltonian is n -degree stable.
- (ii) The property of being traceable is $n - 1$ -degree stable.
- (iii) The property of containing a C_s , ($5 \leq s \leq n$) is $(2n - 1)$ -degree stable.

For other work related to the idea of closure, see [3], [11], [60], [141, 142], [170].

The fourth fundamental result took a different approach. Let $\beta_0(G)$ denote the independence number of G , that is, the size of a maximal independent set of vertices in G .

Theorem 0.4 [47]. If G is a graph with connectivity k such that $\beta_0(G) \leq k$, then G is hamiltonian.

In the following sections, we shall see that each of these results has inspired many others.

1. GENERALIZATIONS OF THE FUNDAMENTALS

Many generalizations of Theorems 0.1-0.4 have been found. Häggkvist and Nicoghossian [86] sharpened Dirac's theorem by incorporating the connectivity of the graph into the degree bound.

Theorem 1.1 [86]. If G is a 2-connected graph of order n , connectivity k and minimum degree $\delta(G) \geq \frac{1}{3}(n + k)$, then G is hamiltonian.



This result itself was recently generalized in [16].

Theorem 1.2 [16]. If G is a 2-connected graph of order n and connectivity k such that $\sigma_3(G) \geq n + k$, then G is hamiltonian.

A natural direction, taken by Bondy [32], was to further increase the number of vertices involved in the independent set.

Theorem 1.3 [32]. If G is a k -connected graph of order $n \geq 3$ such that $\sigma_{k+1}(G) > \frac{1}{2}(k+1)(n-1)$, then G is hamiltonian.

Degree sum conditions like those of Theorems 0.2 and 1.3 do have a major shortcoming, however; they apply to very few graphs. Thus, it is natural to consider variations on such conditions, with the hope that these variations will be more applicable.

Along these same lines, Bondy and Fan [34] provided an Ore-type result for finding a *dominating cycle*, that is, a cycle that is incident to every edge of the graph. Harary and Nash-Williams [89] showed that the existence of a dominating cycle in G is essentially equivalent to the existence of a hamiltonian cycle in the line graph of G , denoted $L(G)$.

Theorem 1.4 [34]. Let G be a k -connected ($k \geq 2$) graph of order n . If any $k+1$ independent vertices x_i ($0 \leq i \leq k$) with $N(x_i) \cap N(x_j) = \emptyset$ ($0 \leq i \neq j \leq k$) satisfy $\sigma_{k+1}(G) \geq n - 2k$, then G contains a dominating cycle.

This result has the immediate corollary that if G is k -connected with $\delta(G) \geq (n - 2k)/(k + 1)$, then G has a dominating cycle. This proves a conjecture of Clark, Colburn, and Erdős [48]. Fraisse [73] had independently proved this conjecture; however, his result is slightly weaker than that of Bondy and Fan.

Bondy [32] also gave a sufficient condition for G to contain a cycle C with the property that $G - V(C)$ contains no clique K_k . When $k = 1$, this result corresponds to Ore's theorem. Veldman [162] further generalized this idea. A cycle C is said to be D_λ -cyclic if and only if every connected subgraph of order λ has at least one vertex in common with C . This idea also generalizes the idea of a dominating cycle. Veldman [162] generalized Theorem 1.1 as well as others to D_λ -cycles.

Another very interesting approach was introduced by Fan [59]. He showed that we need not consider "all pairs of nonadjacent vertices," but only a particular subset of these pairs.

Theorem 1.5 [59]. If G is a 2-connected graph of order n such that

$$\min\{\max(\deg u, \deg v) \mid \text{dist}(u, v) = 2\} \geq \frac{n}{2},$$

then G is hamiltonian.

Fan's theorem is significant generalization of Dirac's theorem. But an entirely new avenue for investigating local structure, along with a search for new adjacency results and bounds," but the set of vertices in question will be: Can an even increasing the number of graphs for that this idea can be used in and that in incorporating more vertices can be useful.

Theorem 1.5 was strengthened to imply the graph is

Recently, a new "generalization unions has proven to be useful set S of vertices. The degree

where $N(v) = \{x \in V(G) \mid x$
 S is chosen to have some relaxation further generalization 1970s and offers a wide variety

The first use of the generalization of Dirac's theorem

Theorem 1.6 [62]. If G is a $(2n - 1)/3$ for each $S = \{x$, then G is hamiltonian.

Fraisse [74] extended this

Theorem 1.7 [74]. Let G there exists some $t \leq k$, such with cardinality t we have d

Very recently, Lindquester tion to vertices at distance t providing an improvement

Theorem 1.8 [110]. If G $\deg(S) \geq (2n - 1)/3$ for every then G is hamiltonian.

Fan's theorem is significant for several reasons. First it is a direct generalization of Dirac's theorem. But more importantly, Fan's theorem opened an entirely new avenue for investigation—one that incorporates some of the local structure, along with a density condition. Now, when attempting to find new adjacency results, one must not only consider the "degree bounds," but the set of vertices for which this bound applies. A natural question will be: Can an even sparser set of vertices be used (thus expanding the number of graphs for which the result will apply)? We shall see later that this idea can be used in conjunction with other adjacency conditions and that incorporating more of the structure beyond the neighborhood of a vertex can be useful.

Theorem 1.5 was strengthened in [22], where the same conditions were shown to imply the graph is pancyclic, with a few minor exceptions.

Recently, a new "generalized degree" approach based upon neighborhood unions has proven to be useful. This idea is based on the adjacencies of a set S of vertices. The degree of a set S is defined to be

$$\text{deg}(S) = \left| \bigcup_{v \in S} N(v) \right|,$$

where $N(v) = \{x \in V(G) \mid xv \in E(G)\}$ is the neighborhood of v . Typically, S is chosen to have some property \hat{P} (for example, independence). This relaxation further generalizes the approach taken in the 1960s and early 1970s and offers a wide variety of uses.

The first use of the generalized degree condition was to provide another generalization of Dirac's theorem.

Theorem 1.6 [62]. If G is a 2-connected graph of order n such that $\text{deg}(S) \geq (2n - 1)/3$ for each $S = \{x, y\}$ where x and y are independent vertices of G , then G is hamiltonian.

Fraisse [74] extended this result to larger independent sets of vertices.

Theorem 1.7 [74]. Let G be a k -connected graph of order n . Suppose there exists some $t \leq k$, such that for every independent set S of vertices with cardinality t we have $\text{deg}(S) \geq t(n - 1)/(t + 1)$, then G is hamiltonian.

Very recently, Lindquester [110] was able to show that a Fan-type restriction to vertices at distance two could also be used with generalized degrees, providing an improvement to Theorem 1.6.

Theorem 1.8 [110]. If G is a 2-connected graph of order n satisfying $\text{deg}(S) \geq (2n - 1)/3$ for every set $S = \{x, y\}$ of vertices at distance 2 in G , then G is hamiltonian.

Independent sets are not the only ones that have been useful in conjunction with generalized degrees. The collection of all pairs of vertices (or all t -sets of vertices) provides yet another generalization of Dirac's Theorem; one with a more combinatorial flavor.

Theorem 1.9 [61]. If G is a 2-connected graph of sufficiently large order n such that $\deg(S) \geq n/2$ for every set S of two distinct vertices of G , then G is hamiltonian.

A similar result holds for sets of more than two vertices (see [61]); however, at this time the best known lower bound is $n/2 + c(k)$, where $c(k)$ is a constant that depends upon k , the number of vertices in the set.

A direct generalization of Ore's theorem was provided in [81]. We say that G satisfies $IC_t(G) \geq k$ if for any set of t independent vertices x_1, \dots, x_t , $|\bigcap_{i=1}^t N(x_i)| \geq k$. A (t_1, t_2) -pair A, B is a pair of sets of independent vertices satisfying $|A| = t_1$, $|B| = t_2$ and $A \cap B = \emptyset$.

Theorem 1.10 [81]. Let G be a graph of order n satisfying $IC_2(G) \geq k$, ($k \geq 2$). Further, let t_1 and t_2 be positive integers satisfying $2 \leq t_1 + t_2 \leq k + 1$. If for every (t_1, t_2) -pair A and B , $\deg(A) + \deg(B) \geq n$, then G is hamiltonian.

Many other results have been discovered in the last few years using generalized degree (neighborhood union) conditions. For a survey of such results see [108].

By varying the typical degree sum approach to that of adjacent vertices rather than nonadjacent vertices, Brualdi and Shanny [42] obtained a hamiltonian result about the *line graph*, $L(G)$, of the given graph.

Theorem 1.11 [42]. If G is a graph of order $n \geq 4$ such that for any edge uv in G , $\deg u + \deg v \geq n$, then G contains a dominating circuit, hence $L(G)$ is hamiltonian.

Veldman [161] further developed this idea. His work can be viewed as yet another form of generalized degree. We follow his notation here. Call two subgraphs H_1 and H_2 of G *close* in G , if they are disjoint and there is an edge of G joining a vertex in H_1 and a vertex of H_2 . If H_1 and H_2 are disjoint, but not close, then they are said to be *remote*. The degree of an edge e of G is the number of vertices of G close to e when e is viewed as a subgraph of order two. We denote the edge degree as $\deg(e)$. Clearly, this is nearly the generalized degree of an adjacent pair of vertices.

Theorem 1.12 [161]. Let G be a k -connected graph ($k \geq 2$) such that for every $k + 1$ mutually remote edges e_0, e_1, \dots, e_k of G , $\sum_{i=0}^k \deg(e_i) > \frac{1}{2}k(n - k)$, then G contains a dominating cycle.

Veldman further conjectures that G is hamiltonian if $\deg(w_i) \geq (n - 2)$. In [20], this work was extended. Veldman also used this approach in [161].

Ainouche and Christofides [2] used this approach under conditions on degrees to obtain a hamiltonian graph (V, E) , with $W \subseteq V$, let

$$\deg w_i \leq d_i$$

be the degrees in G of the vertices w_i . A vertex w_i is "good" if $\deg w_i > i$ for every i . Christofides [2] obtained the following result.

Theorem 1.13 [2]. Let G be a graph of order n . If $\deg x + \deg y \geq n - 2$ for every pair of vertices $x, y \in V - W$, then G is hamiltonian.

Ainouche and Christofides [2] also obtained a result on hamiltonian graphs failing to satisfy Dirac's condition.

Theorem 1.14 [169]. Every 2-regular graph with $\delta(G) \geq (n - 2)/2$ at most $2m$ is hamiltonian. Jackson [169] obtained a k -regular graph with $n \leq 3k$ and $\delta(G) \geq k/2$ is hamiltonian. Jackson's improvement, given in [169].

Theorem 1.14 [169]. Every 2-regular graph with $n \leq 3k + 1$, except for $(3, 2)$ and $(4, 2)$, is hamiltonian.

Recently, a short proof of Jackson's result was given by Kouider [37].

Asratyan and Khachatryan [169] obtained a necessary condition that is reminiscent of Dirac's condition. Let $G_2(x)$ denote the subgraph induced by most 2 from x .

Theorem 1.15 [10]. Let G be a graph of order n . If $\deg x \leq (n - 1)/2$ and y is a vertex of $G_2(x)$, then G is hamiltonian.

Another Ore-type result is given by the following theorem.

Theorem 1.16 [87]. Let G be a graph of order n containing a cycle $C: x_1, x_2, \dots, x_n, x_1$. Suppose G is either pancyclic, bipartite, or a cycle with a chord.

Moreover, if case 3 occurs, then G contains a hamiltonian path. The formation on the local structure of G is crucial.

been useful in conjunction of Dirac's Theorem;

sufficiently large order n at vertices of G , then G

vertices (see [61]); how-
 $\frac{1}{2} + c(k)$, where $c(k)$ is
 tices in the set.
 ovided in [81]. We say
 dent vertices x_1, \dots, x_i ,
 of independent vertices

satisfying $IC_2(G) \geq k$,
 ers satisfying $2 \leq t_1 +$
 $t_2 + \deg(B) \geq n$, then G

the last few years using
 is. For a survey of such

that of adjacent vertices
 hanny [42] obtained a
 he given graph.

such that for any edge uv
 minating circuit, hence

ork can be viewed as yet
 notation here. Call two
 disjoint and there is an
 t_2 . If H_1 and H_2 are dis-
 The degree of an edge e
 en e is viewed as a sub-
 s $\deg(e)$. Clearly, this is
 of vertices.

graph ($k \geq 2$) such that
 $\deg(e_k) > \sum_{i=0}^k \deg(e_i)$

Veldman further conjectures that this bound can be improved to $\frac{1}{3}(k + 1)(n - 2)$. In [20], this work was extended to pancyclic line graphs. Veldman also used this approach in [161] in studying D_λ -cycles.

Ainouche and Christofides [2] combined Pósa [135] and Ore [132] type conditions on degrees to obtain interesting new results. In a graph $G = (V, E)$, with $W \subseteq V$, let

$$\deg w_1 \leq \deg w_2 \leq \dots \leq \deg w_{|W|}$$

be the degrees in G of the vertices in W . A subset W of $V(G)$ is termed "good" if $\deg w_i > i$ for every $w_i \in W$. With this in mind, Ainouche and Christofides [2] obtained the following:

Theorem 1.13 [2]. Let G be a graph of order n and W be a good subset of $V(G)$. If $\deg x + \deg y \geq n$ for any two nonadjacent vertices x, y in $V - W$, then G is hamiltonian.

Ainouche and Christofides also obtained descriptions of maximal non-hamiltonian graphs failing to satisfy their condition. Dirac's condition ($\delta(G) \geq (n/2)$) implies that any m -regular graph of order at most $2m$ is hamiltonian. Jackson [95] has shown that every 2-connected, k -regular graph with $n \leq 3k$ vertices has a hamiltonian cycle. The following improvement, given in [169], was originally conjectured by Jackson.

Theorem 1.14 [169]. Every 2-connected k -regular graph G is hamiltonian if its order $n \leq 3k + 1$, except the Petersen graph.

Recently, a short proof of Theorem 1.14 was found by Bondy and Kouider [37].

Asratyan and Khachatryan [10] introduced yet another Ore-type adjacency condition that is reminiscent of Fan's use of vertices at distance two. Let $G_2(x)$ denote the subgraph of G induced by those vertices at distance at most 2 from x .

Theorem 1.15 [10]. Let G be a graph of order n . Suppose that whenever $\deg x \leq (n - 1)/2$ and y is a vertex at distance 2 from x , $\deg x + \deg_{G_2(x)} y \geq |V(G_2(x))|$, then G is hamiltonian.

Another Ore-type result is due to Hakimi and Schmeichel [87].

Theorem 1.16 [87]. Let G be a graph of order $n \geq 3$ with a hamiltonian cycle $C: x_1, x_2, \dots, x_n, x_1$. Suppose that $\deg x_1 + \deg x_n \geq n$. Then G is either pancyclic, bipartite, or missing only an $(n - 1)$ -cycle.

Moreover, if case 3 occurs, they are able to provide a great deal more information on the local structure around the vertices x_1 and x_n on C .

Denote by $\omega(G)$, the number of components of a graph G . Using this parameter, Chvátal [46] introduced the following concept: We say that G is t -tough if for every vertex cut-set S , $\omega(G - S) \leq |S|/t$. Chvátal showed that if G is hamiltonian, then $t \geq 1$. He also conjectured that if G was 2-tough, then G was hamiltonian. Thomassen and others have produced examples of nonhamiltonian graphs with $t > 3/2$, while in [58] it is shown that there are nonhamiltonian graphs with toughness arbitrarily close to two. Molluzzo [126] also studied toughness. Note that recognizing toughness has recently been shown to be an NP-complete problem [17]. This had been a long-standing open problem.

Toughness, when combined with other conditions, can be used to obtain both new results and improvements of existing results. (See also [25] and [99].)

Theorem 1.17 [98]. Let G be a 1-tough graph of order $n \geq 11$ such that $\sigma_2(G) \geq n - 4$. Then G is hamiltonian.

Theorem 1.18 [18]. Let G be a 2-tough graph of order n such that $\sigma_3(G) \geq n$. Then G is hamiltonian.

Further generalizations of Theorem 1.17 can be found in [147] and generalizations of Fan's theorem with regard to toughness can be found in [15]. For a more complete survey of results relating toughness and hamiltonian properties, see [19].

Turning to work related to Theorem 0.4, we find that in [36] it was shown that a 2-connected graph with $\beta_0(G) \leq 2$ is either pancyclic, or one of the graphs C_4 or C_5 . Amar, Fournier, Germa, and Häggkvist [7] showed that if G is k -connected with $\beta_0(G) = k + 1$, then for every maximum length cycle C of G , $G - V(C)$ is complete. More recently, Benhocine and Fouquet [21] considered hamiltonian line graphs in this context.

Theorem 1.19 [21]. If G is a 2-connected graph and $\beta_0(G) \leq k(G) + 1$, then $L(G)$ is pancyclic unless G is one of C_4, C_5, C_6, C_7 , the Petersen graph or the graph of Figure 1.




FIGURE 1

Many results related to Theorem In 1981, Bermond and Thomassen [and many other results on cycles i some subsequent work related to o

If D is a digraph and $S \subset V(D)$ the digraph induced by S , denoted $D[S]$, is β_1 -independent if $D[S]$ contains no 2-cycles. Thus D is β_1 -independent if $D[S]$ contains no 2-cycles. Thus obtained from a graph G by replacing each edge by a directed edge then $\beta_0 = \beta_1 = \beta_2$. Thus, each p analogue of the undirected indepe

Thomassen [154] gave examples with $\beta_2(D) = 2$ and nonhamiltonian and $\beta_0 = 2$. Thus, the Erdős-Chvátal conjecture generalizes to digraphs. The following p

Problem. Determine if for every (smallest) $f_i(m)$ ($i = 0, 1$, or 2) such that D is β_i -independent with $\beta_i(D) \leq m$ is hamiltonian.

Jackson [96] and Jackson and C

Theorem 1.20 [96].

1. Let D be a digraph with $\beta_0(D) \leq 2$ and D is hamiltonian.
2. Let D be a digraph such that D is β_0 -independent and D has symmetric subgraphs. If $k(l) \geq 2$ and $l \geq 3$, then D is hamiltonian.

A digraph is said to be 2-cyclic if it contains a common cycle.

Theorem 1.21 [97]. If D is a k -connected digraph with $\beta_0(D) \leq k + 1$ and D is 2-cyclic, then D is hamiltonian.

1. if $k \geq 2\beta_1(D) - 1$, then D is hamiltonian;
2. if $k \geq 3$, and $\beta_0(D) \leq 2$, then D is hamiltonian;
3. if $k \geq 15$ and $\beta_0(D) \leq 3$, then D is hamiltonian;
4. if $k \geq 1$ and $\beta_0(D) = 1$, then D is hamiltonian for $3 \leq l \leq n$;
5. if $k \geq 3$ and $\beta_2(D) \leq 2$, then D is hamiltonian for $2 \leq l \leq n$.

Jackson and Ordaz [97] also

Problem.

1. Does there exist an integer k such that every k -connected digraph with $\beta_0(D) = 2$ is hamiltonian?

its of a graph G . Using this concept: We say that G is t -tough if $|S|/t \leq |C|$. Chvátal showed that a graph is t -tough if and only if it is t -connected. It is known that if G was 2-tough, then G is 2-connected. Molluzzo and Molluzzo have produced examples of graphs with toughness close to two. Molluzzo and Molluzzo's toughness has recently been studied [17]. This had been a long-

standing problem, can be used to obtain new results. (See also [25])


graphs of order $n \geq 11$ such that

graphs of order n such that

can be found in [147] and generalizations can be found in [15]. The toughness and hamiltonian

we find that in [36] it was shown that 2 is either pancyclic, or one vertex, and Häggkvist [7] showed that, then for every maximum degree d , there are recently, Benhocine and Benhocine in this context.

graph and $\beta_0(G) \leq k(G) + 1$, C_6, C_7 , the Petersen graph



Many results related to Theorems 0.1–0.2 have been found for digraphs. In 1981, Bermond and Thomassen [24] gave an outstanding survey of these and many other results on cycles in digraphs. I shall now briefly mention some subsequent work related to our four fundamental theorems.

If D is a digraph and $S \subset V(D)$, we say that S is β_0 -independent if the digraph induced by S , denoted $D[S]$, contains no arcs; we say that S is β_1 -independent if $D[S]$ contains no cycles; we say that S is β_2 -independent if $D[S]$ contains no 2-cycles. Thus, $\beta_0 \leq \beta_1 \leq \beta_2$ and if D is the digraph obtained from a graph G by replacing each edge of G by a directed 2-cycle, then $\beta_0 = \beta_1 = \beta_2$. Thus, each parameter may be considered a directed analogue of the undirected independence number β_0 .

Thomassen [154] gave examples of nonhamiltonian 2-connected digraphs with $\beta_2(D) = 2$ and nonhamiltonian 3-connected digraphs with $\beta_1 = 3$ and $\beta_0 = 2$. Thus, the Erdős–Chvátal theorem does not completely generalize to digraphs. The following problem was posed by Jackson [96]:

Problem. Determine if for every integer m , there exists an integer (smallest) $f_i(m)$ ($i = 0, 1$, or 2) such that every $f_i(m)$ -connected digraph D with $\beta_i(D) \leq m$ is hamiltonian.

Jackson [96] and Jackson and Ordaz [97] have investigated this problem:

Theorem 1.20 [96].

1. Let D be a digraph with $\beta_2(G) \leq r$. If $k(D) \geq 2^{r+2}!$, then D is hamiltonian.
2. Let D be a digraph such that $V(D)$ can be covered with m complete symmetric subgraphs. If $k(D) > m(m - 1)$, then D is hamiltonian.

A digraph is said to be 2-cyclic if any two of its vertices are contained in a common cycle.

Theorem 1.21 [97]. If D is a k -connected digraph and

1. if $k \geq 2\beta_1(D) - 1$, then D is 2-cyclic;
2. if $k \geq 3$, and $\beta_0(D) \leq 2$, then D is 2-cyclic;
3. if $k \geq 15$ and $\beta_0(D) \leq 3$, then D is 2-cyclic;
4. if $k \geq 1$ and $\beta_0(D) = 1$, then D contains cycles of length l for $3 \leq l \leq n$;
5. if $k \geq 3$ and $\beta_2(D) \leq 2$, then D contains cycles of lengths l , $2 \leq l \leq n$.

Jackson and Ordaz [97] also posed several more problems:

Problem.

1. Does there exist an integer k such that every k -connected digraph D with $\beta_0(D) = 2$ is hamiltonian?

2. Does every k -connected digraph D with $\beta_0(D) \leq k + 1$ have a hamiltonian path?

Conjecture [97]. Given any integer m , there exists a smallest integer $g(m)$ such that every $g(m)$ -connected digraph D with $\beta_0(D) \leq m$ is 2-cyclic.

In a yet unpublished manuscript, Häggkvist (private communication) proved a Dirac type result on digraphs, namely, that there exists a positive constant c such that every digraph of order n with minimum in- and outdegree at least $(\frac{1}{2} - c)n$ is hamiltonian. Further, he showed that there exist infinitely many digraphs with minimum in- and outdegree at least $(\frac{1}{3} + c)n$ that are not hamiltonian. A basic unsolved problem in this area is to find the best result of this type.

In [30], it is shown that for every positive constant ε , every sufficiently large tournament with minimum in- and outdegree at least $(\frac{1}{4} + \varepsilon)n$ contains the k th power of a hamiltonian cycle.

2. RANDOM GRAPHS AND THE USE OF PROBABILITY

In this section we shall see that probabilistic methods are very useful in studying hamiltonian graphs (and many other graph properties). It is not my purpose to introduce the reader to random graph techniques. For those not familiar with these ideas, see [28].

We shall use $Pr(X)$ to denote the probability of event X . If Ω_n is a model of random graphs of order n , we say *almost every* graph in Ω_n has property Q if $Pr(Q) \rightarrow 1$ as $n \rightarrow \infty$. Note that this is equivalent to saying that the proportion of all labeled graphs of order n that have Q tends to 1 as $n \rightarrow \infty$.

These are two fundamental models for defining probability measures on the set of all 2^M subgraphs (here $M = \binom{n}{2}$) of an n vertex complete graph. Both of these models have been extensively studied.

- (The edge density model) Suppose that $0 \leq p \leq 1$. Let $G_{n,p}$ denote a graph on n vertices obtained by inserting any of the M possible edges with probability p .
- (The fixed size model) Suppose that $N = N(n)$ is a prescribed function of n that takes on values in the set of positive integers. Then there are $S = \binom{M}{N}$ different graphs with N edges possible on the vertex set $\{1, 2, \dots, n\}$. We let $G_{n,N}$ denote one of these graphs chosen uniformly at random with probability $1/S$.

The first major advance in this area was achieved independently by Pósa [136] and Korshunov [103], when they proved the following result:

Theorem 2.1 [136,103]. There exists a constant c such that almost every labeled graph on n vertices and at least $cn \log n$ edges is hamiltonian.

It is also clear that if G is a hamiltonian graph with $\delta(G) \geq 2$. Thus, we see that

$Pr(G_{n,M}$ is hamiltonian)

Komlos and Szemerédi [102] and Komlos [104] showed a threshold for $\delta(G) \geq 2$ with the threshold known that

$$Pr(\delta(G_{n,M}) \geq 2) \sim \omega(n) = 2M/n - \log n$$

They showed that this necessary condition is also sufficient, that almost every $G_{n,M}$ and $G_{n,p}$ is hamiltonian.

Theorem 2.3 [102,104]. Suppose

$$p = \frac{1}{n} \{\log n + \omega(n)\}$$

and

$$L(n) = \left[\frac{n}{2} \{\log n + \omega(n)\} \right]$$

Then almost every $G_{n,p}$ is hamiltonian.

In fact, they showed an even more precise result:

Theorem 2.4 [102,104]. Assume p is constructed as follows: the first edge is chosen at random from the remaining edges of the graph with minimum degree 2 is hamiltonian and the resulting graph is hamiltonian almost surely.

Theorem 2.4 provides us with a precise condition for a graph to be hamiltonian: Simply check if the minimum degree is 0 or 1. The number of times a graph is hamiltonian is given by the number of times the minimum degree is 2 or more.

Related investigations were made by Komlos and Szemerédi [102] and Fenner, and Frieze [29] and Frieze and Komlos [104]. Improvements will be discussed in the next section.

Bollobás et al. [29] used the following result to show that almost every graph with minimum degree 2 is hamiltonian.

Theorem 2.5 [29]. Let G be a graph with minimum degree 2 and n vertices. Then almost every such graph is hamiltonian.

It is also clear that if G is a hamiltonian graph, then its minimum degree $\delta(G) \geq 2$. Thus, we see that

$$Pr(G_{n,M} \text{ is hamiltonian}) \leq Pr(\delta(G_{n,M}) \geq 2).$$

Komlos and Szemerédi [102] and Korshunov [104] were the first to link the threshold for $\delta(G) \geq 2$ with the threshold for G being hamiltonian. It was known that

$$Pr(\delta(G_{n,M}) \geq 2) \longrightarrow 1 \text{ if, and only if, } \omega(n) = 2M/n - \log n - \log \log n \longrightarrow \infty.$$

They showed that this necessary condition was also sufficient to ensure that almost every $G_{n,M}$ and $G_{n,p}$ is hamiltonian.

Theorem 2.3 [102, 104]. Suppose $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$, and let

$$p = \frac{1}{n} \{ \log n + \log \log n + \omega(n) \}$$

and

$$L(n) = \left\lfloor \frac{n}{2} \{ \log n + \log \log n + \omega(n) \} \right\rfloor.$$

Then almost every $G_{n,p}$ is hamiltonian and almost every $G_{n,L}$ is hamiltonian.

In fact, they showed an even more direct relationship.

Theorem 2.4 [102, 104]. Assume that a random labeled graph is constructed as follows: the first edge is chosen at random, the second edge is chosen at random from the remaining $\binom{n}{2} - 1$ possibilities, etc., until a graph with minimum degree 2 is formed. Then the probability that the resulting graph is hamiltonian approaches 1 as $n \rightarrow \infty$.

Theorem 2.4 provides us with an "almost sure decision rule" to decide if a graph is hamiltonian: Simply check whether it contains vertices of degree 0 or 1. The number of times we will be wrong is negligible for large n .

Related investigations were made by Shamir [144], Bollobás [27], Bollobás, Fenner, and Frieze [29] and Frieze [75]. The algorithmic aspects of these improvements will be discussed in Section 4.

Bollobás et al. [29] used the following strengthening of Theorem 2.3 due to Komlos and Szemerédi [102] to produce their algorithmic work.

$\leq k + 1$ have a hamil-

s a smallest integer $g(m)$
($D) \leq m$ is 2-cyclic.

private communication)
t there exists a positive
with minimum in- and
, he showed that there
and outdegree at least
l problem in this area is

unt ϵ , every sufficiently
e at least $(\frac{1}{4} + \epsilon)n$ con-

ABILITY

ods are very useful in
h properties). It is not
h techniques. For those

ent X . If Ω_n is a model
aph in Ω_n has property
lent to saying that the
 Q tends to 1 as $n \rightarrow \infty$.
robability measures on
vertex complete graph.

≤ 1 . Let $G_{n,p}$ denote a
of the M possible edges

i) is a prescribed func-
positive integers. Then
s possible on the vertex
ese graphs chosen uni-

ved independently by
he following result:

such that almost every
es is hamiltonian.

Theorem 2.5 [102]. For $L(n) = (n/2)(\log n + \log \log n + c_n)$

$$\lim_{n \rightarrow \infty} \Pr(G_{n,L} \text{ is hamiltonian}) = \begin{cases} 0, & \text{if } c_n \rightarrow -\infty; \\ e^{-e^{-c}}, & \text{if } c_n \rightarrow c; \\ 1, & \text{if } c_n \rightarrow \infty. \end{cases}$$

For $V_n = \{1, 2, \dots, n\}$, let $v \in V_n$ independently make m random (but not necessarily distinct) choices $c(v, i) \in V_n$, $i = 1, 2, \dots, m$. This is done independently for each $v \in V_n$. Then consider the multigraph

$$D(n, m) = (V_n, E(n, m)), \text{ where}$$

$$E(n, m) = \{(v, c(v, i)) \mid v \in V_n, 1 \leq i \leq m, \text{ and } v \neq c(v, i)\}.$$

(That is, we ignore the orientation on the edges $(v, c(v, i))$, but we do not coalesce multiple edges or remove loops. Then with this in mind, Fenner and Frieze [66] accomplished a major step when they verified these graphs are almost always hamiltonian. Their proof was the first example of the "coloring technique" that has proved most useful in this area.

Theorem 2.6 [66]. For $m \geq 23$, $\lim_{n \rightarrow \infty} \Pr(D(n, m) \text{ is hamiltonian}) = 1$.

They further conjecture the naturally anticipated fact that this can be improved to $m \geq 3$. Frieze [75] was able to improve this to $m \geq 10$ as well as improve the time of the algorithm used to produce the cycle (see Section 4 for more details).

Let $R(n, r)$ denote the random regular graph chosen uniformly from the set of r -regular graphs on V_n . Bollobás [26] and Fenner and Frieze [67] independently proved that there is a constant r_0 such that for any $r \geq r_0$,

$$\lim_{n \rightarrow \infty} \Pr(R(n, r) \text{ is hamiltonian}) = 1.$$

In [67], it was shown that $r_0 = 796$, while in [75], this was improved to $r_0 = 85$. Again, Frieze conjectures that the best value actually is $r_0 = 3$.

One might hope that the problem of finding hamiltonian cycles in random bipartite graphs is easier than in $G_{n,p}$. However, this is not the case. Progress was made by Frieze [76]. Here we let $G_{n,n;p}$ denote a random bipartite graph with n vertices in each partite set and probability p that any edge is in $G_{n,n;p}$.

Theorem 2.7 [76]. Let $p = ((\log n + \log \log n + c_n)/n)$. Then the probability that $G_{n,n;p}$ is hamiltonian tends to $e^{-2e^{-c}}$ as $c_n \rightarrow c$.

As with random graphs, the obstacle to be overcome in random bipartite graphs turns out to be the existence of vertices of degree at most 1.

Turning to digraphs, we note the result (especially in view of the fact that the problem (for more details) does not have a solution) was proved by McDiarmid [122, 123] was able to show that a random digraph $D_{n,p}$ is hamiltonian if $G_{n,p}$ is hamiltonian. Using this fact

Theorem 2.8 [122, 123]. If $p =$

$$\Pr(D_{n,p} \text{ is hamiltonian}) =$$

Other interesting results are proved that the probability that a random digraph is hamiltonian is 0.974. They also showed that almost all digraphs are hamiltonian. However, Richmond, Roberson, and others have shown that sometimes hamiltonian cycles are rare.

Theorem 2.9 [137]. Almost every

3. FORBIDDEN SUBGRAPHS

A new approach to the hamiltonian cycle problem in general, began with Alon and Hedetniemi [77]. Before this, the problem was not helpful. Given graphs F_1, F_2 , a graph G is (F_1, F_2) -free if G contains no induced subgraph isomorphic to F_1 or F_2 .

In considering graphs that are forbidden, our attention to a class of graphs is drawn. Thus, we may be able to find results that apply when the graphs are not previously covered by density results that apply when the graphs are forbidden. Central to most forbidden subgraph theory is the complete bipartite graph $K_{1,3}$ (sometimes called a star $K_{1,3}$ (see Figure 2). Some other forbidden subgraphs are shown in Figure 3.

We are now ready to state

Theorem 3.1 [77]. If G is a graph with no induced $K_{1,3}$ subgraph, then G is hamiltonian.

The proof of Theorem 3.1 shows that the only graphs satisfying

$\log \log n + c_n$)

- if $c_n \rightarrow -\infty$;
- if $c_n \rightarrow c$;
- if $c_n \rightarrow \infty$.

ly make m random (but not \dots, m . This is done inde- multigraph

where

i , and $v \neq c(v, i)$).

s ($v, c(v, i)$), but we do not with this in mind, Fenner they verified these graphs s the first example of the d in this area.

\dots, m) is hamiltonian) = 1.

ated fact that this can be ove this to $m \geq 10$ as well roduce the cycle (see Sec-

hosen uniformly from the onner and Frieze [67] inde- that for any $r \geq r_0$,

) = 1.

75], this was improved to alue actually is $r_0 = 3$.

hamiltonian cycles in ran- ever, this is not the case.

$D_{n,n;p}$ denote a random bi- and probability p that any

c_n/n). Then the proba- $c_n \rightarrow c$.

come in random bipartite f degree at most 1.

Turning to digraphs, we note that the analogous problem seems harder, especially in view of the fact that the useful work of Pósa [136] (see Section 4 for more details) does not have directed analogues. But despite this problem, McDiarmid [122, 123] was able to show that the probability that a random digraph $D_{n,p}$ is hamiltonian is not smaller than the probability that $G_{n,p}$ is hamiltonian. Using this fact he deduced the following result:

Theorem 2.8 [122, 123]. If $p = (1/n)(1 + \epsilon)(\log n)$ then

$$Pr(D_{n,p} \text{ is hamiltonian}) \rightarrow \begin{cases} 1, & \text{if } \epsilon > 0; \\ 0, & \text{if } \epsilon < 0. \end{cases}$$

Other interesting results are due to Robinson and Wormald [140], who proved that the probability that a cubic graph is hamiltonian is at least 0.974. They also showed that almost every cubic bipartite graph is hamiltonian. However, Richmond, Robinson, and Wormald [137] showed that at times hamiltonian cycles are rare.

Theorem 2.9 [137]. Almost every cubic planar graph is nonhamiltonian.

3. FORBIDDEN SUBGRAPHS

A new approach to the hamiltonian problem, although not new to graph theory in general, began with a rather innocent observation due to Goodman and Hedetniemi [77]. Before exploring this approach, some terms will be helpful. Given graphs F_1, F_2, \dots, F_k , we say that G is $\{F_1, F_2, \dots, F_k\}$ -free if G contains no induced subgraph isomorphic to any F_i ($1 \leq i \leq k$).

In considering graphs that are free of some set of graphs, we are restricting our attention to a class of graphs defined with specific structural limitations. Thus, we may be able to avoid the pure density-type arguments seen earlier. Our hope, of course, is to find conditions that will work on graphs not previously covered by density results. In fact, what we tend to obtain are results that apply when the graphs are either dense or very sparse.

Central to most forbidden subgraph results to date is the complete bipartite graph $K_{1,3}$ (sometimes called a *claw*) or graphs very closely related to $K_{1,3}$ (see Figure 2). Some other graphs that have proven to be useful are shown in Figure 3.

We are now ready to state Goodman and Hedetniemi's result.

Theorem 3.1 [77]. If G is a 2-connected $\{K_{1,3}, Z_1\}$ -free graph, then G is hamiltonian.

The proof of Theorem 3.1 is very simple and in fact it is easy to show that the only graphs satisfying its hypothesis are complete graphs, complete




FIGURE 2

graphs with a matching removed, or a cycle. Goodman and Hedetniemi pointed out that this seemed to be the first result that actually applied to a cycle.

In 1979, Oberly and Sumner [131] really opened the door to this approach, by relating forbidden subgraphs with another property, local connectivity. We say a graph G is locally connected, if for each vertex x , the subgraph of G induced by $N(x)$ is a connected graph.

Theorem 3.2 [131]. A connected, locally connected, $K_{1,3}$ -free graph of order $n \geq 3$ is hamiltonian.

Further, Oberly and Sumner made the following interesting conjecture:

Conjecture. If G is a connected, locally k -connected, $K_{1,k+2}$ -free graph of order $n \geq 3$, then G is hamiltonian.

The work of Oberly and Sumner spurred further investigations of the same type. Attempts were made to broaden the sets of graphs that were forbidden. See Figures 2 and 3 for some of the graphs that have been used.

Theorem 3.3 [54]. Let G be a graph of order $n \geq 3$ that is $\{K_{1,3}, F\}$ -free. Then,

- (i) if G is connected, then G is traceable;
- (ii) if G is 2-connected, then G is hamiltonian.

This result was followed by other

Theorem 3.4 [79]. If G is a 2-connected graph, then G is pancyclic or G is a cycle.

Since I and A are induced subgraphs, they are also F -free. Thus, the following corollary holds.

Corollary 3.5. Let G be a 2-connected graph.

- (i) If G is I -free, then G is hamiltonian.
- (ii) If G is A -free, then G is hamiltonian.

Zhang [168] considered degree conditions. He showed that if G is a k -connected graph and $\sigma_{k+1}(G) \geq n - k$, then G is hamiltonian.

Broersma and Veldman [41] introduced a graph condition by allowing certain forbidden subgraphs. We say a subgraph H of G

$$(N(u) \cap N(v)) \cap H \neq \emptyset$$

That is, $u, v \in V(H)$ and u and v are adjacent in H . Using this idea, they obtained the following theorem (Theorem 3.1). The vertices a, b, c

Theorem 3.6 [41]. Let G be a 2-connected graph.

- (i) If every induced Z_1 of G satisfies $\phi(a, b_1)$, then G is pancyclic or G is a cycle.
- (ii) If every induced Z_2 of G satisfies $\phi(a, b_1)$, then G is pancyclic or G is a cycle.

The nonhamiltonian $K_{1,3}$ -free graph G does not satisfy every induced Z_2 satisfies $\phi(a, b_1)$. Broersma and Veldman cannot be replaced by "or." Broersma and Veldman's realization of Corollary 3.5(i) using related graphs (see Figure 3) to

Theorem 3.7 [41]. Let G be a 2-connected graph. If every induced subgraph of G isomorphic to Z_1 satisfies $\phi(a, c_1)$ and $\phi(a, c_2)$, then G is hamiltonian.

An immediate corollary of Theorem 3.7 is

Corollary 3.8 [78]. If G is a 2-connected graph with $\delta(G) \geq \frac{n}{2}$, then G is hamiltonian.

This result was followed by other extensions of Theorem 3.1.

Theorem 3.4 [79]. If G is a 2-connected $\{K_{1,3}, Z_2\}$ -free graph, then either G is pancyclic or G is a cycle.

Since I and A are induced subgraphs of F , every I -free or A -free graph is also F -free. Thus, the following corollary of Theorem 3.3 is obtained:

Corollary 3.5. Let G be a 2-connected $K_{1,3}$ -free graph.

- (i) If G is I -free, then G is hamiltonian.
- (ii) If G is A -free, then G is hamiltonian.

Zhang [168] considered degree sums in claw free graphs. In particular, he showed that if G is a k -connected, $K_{1,3}$ -free graph of order n such that $\sigma_{k+1}(G) \geq n - k$, then G is hamiltonian.

Broersma and Veldman [41] introduced a relaxation of the forbidden subgraph condition by allowing certain of the forbidden graphs to exist, provided their adjacencies outside their own vertex set are of the "proper type." We say a subgraph H of G satisfies property $\phi(u, v)$ if

$$(N(u) \cap N(v)) - V(H) \neq \emptyset.$$

That is, $u, v \in V(H)$ and u and v have a common neighbor in G outside of H . Using this idea, they obtained generalizations to several results, including Theorem 3.1. The vertices a, b_1 , and b_2 are as in Figure 3.

Theorem 3.6 [41]. Let G be a 2-connected $K_{1,3}$ -free graph.


- (i) If every induced Z_1 of G satisfies $\phi(a, b_1)$ or $\phi(a, b_2)$, then either G is pancyclic or G is a cycle.
- (ii) If every induced Z_2 of G satisfies $\phi(a_1, b_1)$ or $\phi(a_1, b_2)$, then either G is pancyclic or G is a cycle.

The nonhamiltonian $K_{1,3}$ -free graph of Figure 4 has the property that every induced Z_2 satisfies $\phi(a_1, b_1)$ or $\phi(a_1, b_2)$; hence, in Theorem 3.6, "and" cannot be replaced by "or." Broersma and Veldman also obtained a generalization of Corollary 3.5(i) using these ideas. They also used some other related graphs (see Figure 3) to obtain the following result:

Theorem 3.7 [41]. Let G be a 2-connected $K_{1,3}$ -free graph. If every induced subgraph of G isomorphic to P_7 or P_7^+ satisfies $\phi(a, b_1)$ or $\phi(a, b_2)$ or $(\phi(a, c_1) \text{ and } \phi(a, c_2))$, then G is hamiltonian.

An immediate corollary of Theorem 3.7 was originally obtained in [78].

Corollary 3.8 [78]. If G is a 2-connected $K_{1,3}$ -free graph of diameter at most 2, then G is hamiltonian.



Goodman and Hedetniemi
ult that actually applied to
ened the door to this ap-
rother property, local cond,
d, if for each vertex x , the
raph.
ected, $K_{1,3}$ -free graph of
ng interesting conjecture:
nected, $K_{1,k+2}$ -free graph
her investigations of the
s of graphs that were for-
is that have been used.
 ≥ 3 that is $\{K_{1,3}, F\}$ -free.




FIGURE 3

Broersma and Veldman [41] conjectured the following generalization of Corollary 3.5(ii) and Theorem 3.3.

Conjecture.

1. Let G be a 2-connected $K_{1,3}$ -free graph. If every induced A of G satisfies $\phi(a_1, a_2)$, then G is hamiltonian.
2. Let G be a 2-connected $K_{1,3}$ -free graph. If every induced F of G satisfies $(\phi(a_1, a_2)$ and $\phi(a_1, a_3))$ or $(\phi(a_1, a_2)$ and $\phi(a_2, a_3))$ or $(\phi(a_1, a_3)$ and $\phi(a_2, a_3))$, then G is hamiltonian.




FIGURE 4

Recently, a different relaxation in which they showed that if then it is hamiltonian.

Theorem 3.9 [69]. Let G be a graph with minimum degree δ . If $\delta \geq (n-1)/3$ and v , the number of induced $K_{1,3}$ and v is less than $\delta - 1$, then

In [70], Flandrin and Li showed

$$\sigma_3(G) \geq \frac{4n}{3}$$

then G is hamiltonian. This result is related to $|N(w)|$ in [68].

Matthews and Sumner [120] obtained a result for $K_{1,3}$ -free graphs.

Theorem 3.10 [121]. Let G be a graph with $\delta(G) \geq (n-2)/3$, then G is hamiltonian.

Matthews and Sumner also

Conjecture [120]. If G is a graph with $\delta(G) \geq (n-2)/3$ and G is a 4-regular graph, then G is hamiltonian.


It is interesting to note that this conjecture is true when we have a reasonable number of vertices.

Theorem 3.11 [63]. If G is a graph with $\delta(G) \geq (n-2)/3$ and $S = \{x, y\}$, where x and y are adjacent vertices such that $\deg S > (2n-2)/3$, then G is hamiltonian.

Conjecture [63]. If G is a graph with $\delta(G) \geq (n-2)/3$ and S is a set of vertices such that $\deg S > (2n-5)/3$, where S is hamiltonian.

Another problem in this area is the result of Fleischner [71], showing that a graph is hamiltonian. (Recently, Rihaczek [72] gave a counterexample to the typical example that is used in Fleischner's theorem is the claw (see Figure 5), which is not hamiltonian.

In [80], it was conjectured that a graph must be hamiltonian. This conjecture is still open.



Recently, a different relaxation has been explored by Flandrin and Li [69] in which they showed that if a graph does not contain "too many" claws, then it is hamiltonian.

Theorem 3.9 [69]. Let G be a 2-connected graph of order $n \geq 16$ and minimum degree δ . If $\delta \geq (n/3)$ and if for any two nonadjacent vertices u and v , the number of induced subgraphs isomorphic to $K_{1,3}$ containing u and v is less than $\delta - 1$, then G is hamiltonian.

In [70], Flandrin and Li showed that if G is 2-connected and

$$\sigma_3(G) \geq \frac{4n}{3} + |N(u) \cap N(v) \cap N(w)|,$$

then G is hamiltonian. This bound was reduced to $n + |N(u) \cap N(v) \cap N(w)|$ in [68].

Matthews and Sumner [120, 121] studied hamiltonian properties of graphs obtained from $K_{1,3}$ -free graphs.

Theorem 3.10 [121]. Let G be a 2-connected, $K_{1,3}$ -free graph with $\delta(G) \geq (n - 2)/3$, then G is hamiltonian.

Matthews and Sumner also made the following conjecture:

Conjecture [120]. If G is a 4-connected $K_{1,3}$ -free graph, then G is hamiltonian.

It is interesting to note that we can reduce the connectivity from 4 to 2, when we have a reasonable neighborhood union condition present.

Theorem 3.11 [63]. If G is a 2-connected $K_{1,3}$ -free graph of order $p \geq 14$ and $S = \{x, y\}$, where x and y are nonadjacent vertices of G , and for each such S , $\text{deg } S > (2n - 2)/3$, then G is pancyclic,

Conjecture [63]. If G is a 3-connected $K_{1,3}$ -free graph of order n such that $\text{deg } S > (2n - 5)/3$, where S is any set of two nonadjacent vertices, then G is hamiltonian.


Another problem in this area arose from consideration of the famous result of Fleischner [71], showing that the square of any 2-connected graph is hamiltonian. (Recently, Riha [138] has obtained a short proof of this result). The typical example that shows that the connectivity cannot be lowered in Fleischner's theorem is provided by $S(K_{1,3})$, the subdivision graph of the claw (see Figure 5), whose square is not hamiltonian.

In [80], it was conjectured that the square of any connected $S(K_{1,3})$ -free graph must be hamiltonian. This conjecture was verified by Hendry and

Following generalization of

every induced A of G satis-

every induced F of G satis-
d $\phi(a_2, a_3)$ or $\phi(a_1, a_3)$






FIGURE 5

Vogler [91]. We conclude this section with the following powerful result of Fleischner [72] concerning the square of a graph:

Theorem 3.12 [72]. For a connected graph G ,

- (i) G^2 is hamiltonian if and only if G^2 is vertex pancyclic, and
- (ii) G^2 is hamiltonian-connected if and only if G^2 is panconnected.

4. ALGORITHMS

Despite the fact that the hamiltonian problem is NP-complete, algorithms of a probabilistic nature and algorithms for special classes of graphs have been developed. As was mentioned in Section 2, Pósa [136] was the first to suggest an algorithm that converges almost surely for a graph of order n and size $cn \log n$, $c \geq 3$. The ideas behind his theoretic work suggested a probabilistic algorithm for determining the existence of a hamiltonian cycle. Tests of this algorithm were first performed by McGregor (see [100]) on graphs of order up to 500 and by Thompson and Singhal [156] on graphs of order up to 1000. The ideas behind Pósa's work have been refined in [29] and [75] to obtain improvements in time complexity. Here we naturally only consider graphs with minimum degree at least 2.

The fundamental idea behind Pósa's algorithm is a path transformation operation often called a *rotation*. It works as follows: Given a path $P = v_1, v_2, \dots, v_k$ and an additional edge $e = v_k v_i$ ($1 \leq i \leq k - 2$), we can create a new path, also of length $k - 1$, by deleting the edge $v_i v_{i+1}$ and inserting the edge e . Thus, define the path operation ROTATE (P, e) as

$$\text{ROTATE } (P, e) = v_1, v_2, \dots, v_i, v_k, v_{k-1}, \dots, v_{i+1}.$$

The operation, ROTATE produces a new path with v_1 as its initial vertex and v_{i+1} as its end vertex.

Pósa's algorithm begins by selecting a vertex x_0 and trying to extend this trivial path, call it P , by including any unused neighbor of the end vertex (namely, x_0) of this path. At first this extension adds some neighbor, say x_1 to P . We now repeat this step from x_1 and continue extending P from the nonfixed end vertex until we can no longer extend the path. At this point,

either we have a hamiltonian path or a nonfixed end vertex of the path. If we exist an edge $e = v_k v_i$ ($1 \leq i \leq k - 2$), we can rotate (P, e) to obtain a new path, say P' . If we are unable to extend P' , we continue this process until a hamiltonian path is found. The number of rotations exceeds some specific bound, called the *extension-rotation* algorithm.

Other early algorithms were developed by Bollobás et al. [144]. In 1984, Bollobás et al. [144] found finding hamiltonian paths. Their algorithm had time complexity $O(n^{4+\epsilon})$. It is a probabilistic technique. Recently, Frieze [75] used Pósa's techniques can be used to find a hamiltonian path HAM 1, which satisfies

$$\lim_{n \rightarrow \infty} Pr(\text{HAM 1 finds a path}) = 1$$

Further, Luczak and Frieze (see [144]) used Frieze [75] algorithm has shown that HAM 2, which satisfies

$$\lim_{n \rightarrow \infty} Pr(\text{HAM 2 finds a path}) = 1$$

for any constant $r \geq 85$.

Another recent development used an edge coloring based algorithm to construct a hamiltonian path from a graph with minimum degree at least 2.

Theorem 4.1 [83]. There is a probabilistic algorithm: For any fixed probability p and any constant c , $Pr(\text{HAM } 2 \text{ finds a path}) = (cn)/p + o(n)$.

Other special case algorithms exist for graphs with minimum degree at least 2.

5. MULTIPLE HAMILTONIAN PATHS

In trying to construct hamiltonian paths, the transformation from a nonfixed end vertex to a fixed end vertex often many different spanning trees. We count the number of distinct cycles. We wish to show the existence of a hamiltonian path. We consider both of these questions.

We begin with results on edge coloring. The first such results is due to Nešetřil [144] of graphs.

either we have a hamiltonian path and we stop, or we ROTATE from the nonfixed end vertex of the path. Since $\delta(G) \geq 2$, we see that there must exist an edge $e = v_k v_i$ ($1 \leq i \leq k$) and hence we can perform ROTATE (P, e) to obtain a new path, say P' . We now try to extend this new path, rotating when we are unable to extend the (nonhamiltonian) path. We continue this process until a hamiltonian path is found or until the number of rotations exceeds some specified limit. This technique has come to be called the *extension-rotation* approach.

Other early algorithms were due to Angluin and Valiant [8] and Shamir [144]. In 1984, Bollobás et al. [29] developed an improved algorithm for finding hamiltonian paths. Their algorithm almost always succeeded and had time complexity $O(n^{4+\epsilon})$. It was still based on the extension-rotation technique. Recently, Frieze [75] has shown that a careful modification of Pósa's techniques can be used to produce a $O(n^3 \log n)$ time algorithm HAM 1, which satisfies

$$\lim_{n \rightarrow \infty} Pr(\text{HAM 1 finds a hamiltonian cycle in } D(n, 10)) = 1.$$

Further, Luczak and Frieze (see [75]) have reduced the 10 above to 5.

Frieze [75] also has shown that there is an $O(n^3 \log n)$ time algorithm HAM 2, which satisfies

$$\lim_{n \rightarrow \infty} Pr(\text{HAM 2 finds a hamiltonian cycle in } R(n, r)) = 1,$$

for any constant $r \geq 85$.

Another recent development is due to Gurevich and Shelah [83]. They used an edge coloring based algorithm (called HPA) to almost always construct a hamiltonian path from a fixed initial vertex to a fixed final vertex.

Theorem 4.83. There is a positive real number c satisfying the following: For any fixed probability p , the expected run time of HPA on $G_{n,p}$ is $(cn)/p + o(n)$.

Other special case algorithms can be found in [1], [9], [141], and [143].

5. MULTIPLE HAMILTONIAN CYCLES

In trying to construct hamiltonian graphs, it is common to notice that in the transformation from a nonhamiltonian graph to a hamiltonian graph, often many different spanning cycles are created. Thus, at times we wish to count the number of distinct cycles that are present and at other times we wish to show the existence of several edge-disjoint cycles. We shall now consider both of these questions.

We begin with results on edge-disjoint hamiltonian cycles. One of the first such results is due to Nebesky and Wisztova [129] and concerns powers of graphs.

owing powerful result of

pancyclic, and \mathcal{F}^2 is panconnected.

NP-complete, algorithms
 classes of graphs have
 ósa [136] was the first to
 for a graph of order n
 theoretic work suggested a
 e of a hamiltonian cycle.
 McGregor (see [100]) on
 inghal [156] on graphs of
 ave been refined in [29]
 xity. Here we naturally
 st 2.

is a path transformation
 lows: Given a path $P =$
 $1 \leq i \leq k - 2$, we can
 g the edge $v_i v_{i+1}$ and in-
 n ROTATE (P, e) as

$-1, \dots, v_{i+1}$.

th v_1 as its initial vertex

and trying to extend this
 ighbor of the end vertex
 lds some neighbor, say x_1
 ue extending P from the
 l the path. At this point,

Theorem 5.1 [129]. If G is a connected graph of order at least $n \geq 6$, then there exists a hamiltonian cycle C of G^3 and a hamiltonian cycle C_1 of G^5 such that C and C_1 are edge-disjoint.

This result strengthens the well-known results that G^3 is hamiltonian and if $n \geq 5$, that G^5 has a 4-factor.

Other density conditions have been developed along the lines we investigated in Section 1. Nash-Williams [128] generalized Dirac's theorem to obtain a result on multiple edge-disjoint hamiltonian cycles.

Theorem 5.2 [129]. If G is a graph of order n such that $\delta(G) \geq n/2$, then G contains $\lfloor (5(n + a_n + 10))/224 \rfloor$ edge-disjoint hamiltonian cycles, where

$$a_n = \begin{cases} 0, & \text{if } n \text{ is even;} \\ 1, & \text{otherwise.} \end{cases}$$

Jackson [94] investigated multiple hamiltonian cycles in regular graphs.

Theorem 5.3 [94]. If G is a k -regular graph of order n ($n \geq 14$) and $k \geq (n - 1)/2$, then G contains $(3k - n + 1)/6$ edge-disjoint hamiltonian cycles.

Note that Jackson's Theorem provides a strengthening of Theorem 5.2 in the case of regular graphs. Jackson also conjectured that if G is a k -regular graph on n vertices, where $k \geq (n - 1)/2$, then G contains $k/2$ edge-disjoint hamiltonian cycles. That this conjecture cannot be extended to small k ($k = 4, 5$) has been shown by Zaks [165]. He demonstrated an infinite family of 4-regular, 4-connected graphs in which any two hamiltonian cycles shared at least $\frac{1}{16}$ of their edges and he demonstrated a family of 5-regular, 5-connected planar graphs without two edge-disjoint hamiltonian cycles. Such a family of 5-regular graphs was also found by Owens [133]. Owens [133] also showed the existence for every $r \geq 3$, and every $k, 0 \leq k \leq n/2$, of an r -regular, r -connected graph that contains k edge-disjoint hamiltonian cycles, but not $k + 1$ edge disjoint hamiltonian cycles.

Faudree, Rousseau, and Schelp [65] developed a degree sum condition implying the existence of multiple hamiltonian cycles and in so doing produced another generalization of Ore's theorem.

Theorem 5.4 [65]. Let G be a graph of order $n \geq 3$ and k be a positive integer. If the sum of the degrees of any pair of nonadjacent vertices is at least $n + 2k - 2$, then for n sufficiently large ($n \geq 60k^2$ will suffice), G has k edge-disjoint hamiltonian cycles.

They further conjectured that the degree sum condition could be decreased to " $\geq n$," if an additional minimum degree condition was imposed. It should be noted that at the same time, Li and Zhu [109] independently proved the following:

Theorem 5.5 [109]. Let G be a graph of order n such that $\deg x + \deg y \geq n$ for any pair of nonadjacent vertices x, y . Then G contains at least two edge-disjoint hamiltonian cycles.

Faudree et al. [65] were also able to obtain a result for the $k = 1$ case below) based on the following:

Theorem 5.6 [65]. Let G be a graph of order n and size $\binom{n-1}{2} + 2k$.

1. If $n \geq 6k$, then G has k edge-disjoint hamiltonian cycles.
2. If $n \geq 6k^2$, then G has k edge-disjoint hamiltonian cycles in the range 3 to n .

The generalized degree condition obtained by Faudree et al. [65] can be used to obtain a result on multiple edge-disjoint hamiltonian cycles. Additional conditions were not required. The minimum degree condition for a nontrivial graph is the minimum degree condition. This result in a disconnected graph.

Theorem 5.7 [64]. Let G be a graph of order n and constant $c = c(k)$ such that $\delta(G) \geq c$ satisfying

1. $|N(u) \cup N(v)| \geq ((2n + 1)/2) - c$,
2. $\delta(G) \geq 4k + 1$,
3. $k_1(G) \geq 2k$, and
4. $k_1(G - v) \geq k$ for every vertex v .

then G contains k edge-disjoint hamiltonian cycles.

Any result that supplies sufficient conditions for the existence of multiple edge-disjoint hamiltonian cycles like condition 1 must have been proved in [64]. Conditions 2 and 4 are known to be sharp.

A corresponding result using the degree sum condition for pairs of vertices would be interesting. Also, extensions of Theorem 5.4 would be desirable.

Bondy and Häggkvist [35] obtained a result of Grinberg [82].

Theorem 5.8 [35]. Let G be a graph of order n and size $\binom{n-1}{2} + 2k$. Then G contains k edge-disjoint hamiltonian cycles. Also, the sets of faces of G into

Theorem 5.5 [109]. Let G be a graph of order $n \geq 20$ and let $\delta(G) \geq 5$. If $\deg x + \deg y \geq n$ for any pair of nonadjacent vertices x and y , then G contains at least two edge-disjoint hamiltonian cycles.

Faudree et al. [65] were also able to generalize another of Ore's results (the $k = 1$ case below) based on the size of the graph.

Theorem 5.6 [65]. Let k be a positive integer and G a graph of order n and size $\binom{n-1}{2} + 2k$.

1. If $n \geq 6k$, then G has k edge-disjoint hamiltonian cycles.
2. If $n \geq 6k^2$, then G has k edge-disjoint cycles of length l , for any integer l in the range 3 to n .

The generalized degree condition discussed earlier has also been used to obtain a result on multiple edge-disjoint cycles. In order to do this, several additional conditions were necessary. The edge-connectivity, $k_1(G)$, of a nontrivial graph is the minimum number of edges whose removal from G results in a disconnected graph.

Theorem 5.7 [64]. Let k be a fixed positive integer. Then there is a constant $c = c(k)$ such that if G is a graph of sufficiently large order n satisfying

1. $|N(u) \cup N(v)| \geq ((2n + c)/3)$ for each pair u, v of nonadjacent vertices,
2. $\delta(G) \geq 4k + 1$,
3. $k_1(G) \geq 2k$, and
4. $k_1(G - v) \geq k$ for every vertex v ,

then G contains k edge-disjoint hamiltonian cycles.

Any result that supplies sufficient conditions for a graph G to contain k edge-disjoint hamiltonian cycles and is based on a generalized degree condition like condition 1 must have these types of added restrictions. Examples to show this are provided in [64]. However, at this time, only conditions 3 and 4 are known to be sharp.

A corresponding result using all pairs of vertices rather than nonadjacent pairs of vertices would be interesting, but at this time remains unknown. Also, extensions of Theorem 5.7 to the case of more than two vertices would be desirable.

Bondy and Häggkvist [35] developed a generalization of the well-known result of Grinberg [82].

Theorem 5.8 [35]. Let G be a 4-regular plane graph that is decomposable into edge-disjoint hamiltonian cycles C and D . Denote by F_{11}, F_{12}, F_{21} , and F_{22} the sets of faces of G interior to both C and D , interior to C but not D ,

interior to D but exterior to C and exterior to both C and D , respectively. Then

$$g(F_{11}) = g(F_{22}) \quad \text{and} \quad g(F_{12}) = g(F_{21})$$

where $g: 2^F \rightarrow N$ defined by $g(X) = \sum_{f \in X} (d(f) - 2)$ where $d(f)$ is the number of edges in the boundary of f .

Note that Zaks [166] has another generalization of Grinberg's theorem.

The question of counting the number of hamiltonian cycles has been considered in several papers. Sheehan and Wright [146] counted hamiltonian cycles in dense graphs.

Theorem 5.9 [146]. Let G be an (n, q) -graph with $\Delta(G) = \beta$, let $H(G)$ be the number of hamiltonian cycles in G , and let $M = ((n-1)!)/2$ be the number of hamiltonian cycles in K_n . Then, if

$$\frac{q}{n} \longrightarrow a < \infty \quad \text{as} \quad n \longrightarrow \infty \quad \text{and} \quad \beta = o(n), \quad \text{then}$$

$$\frac{H(G)}{M} \longrightarrow e^{-2a} \quad \text{as} \quad n \longrightarrow \infty.$$

Sheehan [145] also studied graphs with exactly one hamiltonian cycle.

Theorem 5.10 [145]. Let G be a graph of order n containing exactly one hamiltonian cycle. Then the maximum number of edges in G is $(n^2/4) + 1$.

As usual, special classes of graphs also provide us with a chance to say more.

Theorem 5.11 [88].

1. For all $n \geq 12$, there exists a maximal planar graph of order n with exactly four hamiltonian cycles.
2. Every 4-connected maximal planar graph on n vertices contains at least $n/\log_2 n$ hamiltonian cycles.

A. Thomason [153] provided the answer to several interesting problems. Smith (see [159]) proved that in a cubic graph, the number of hamiltonian cycles containing a given edge is even. Thomason [153] proved that if all vertices of G , with the possible exception of two (say u and v), have odd degree, then the number of hamiltonian paths from u to v is even. Thomason also generalized in several ways the result of Kotzig (see [40]) that in a bipartite cubic graph, the total number of hamiltonian cycles is even.

Sloane [148] asked if the existence of hamiltonian cycles in G implied the existence of hamiltonian paths. He answered this positively.

Theorem 5.12 [153]. In a 4-regular graph, the number of pairs of edge-disjoint hamiltonian cycles is even.

Nincak [130] proved that if a graph contains at least k hamiltonian cycles, then G contains at least k edge-disjoint hamiltonian cycles. [153] showed the following:

Theorem 5.13 [153]. If a $2k$ -regular graph can be decomposed into k edge-disjoint hamiltonian cycles, then

1. each edge of G is in at least k cycles;
2. G has at least $k(3k-2)$ edges;
3. G has at least $(3k-2)k$ vertices.

Tomescu [157] considered the following problem:

Theorem 5.14 [157]. Let G be a $2m$ -regular graph ($mk \equiv 0 \pmod{2}$).

1. If $k \geq 1$ and $m \geq 3k$, then G has at least $(m-1)(m-2)\cdots(m-k)$ edge-disjoint hamiltonian cycles.
2. The graph G has at least k edge-disjoint hamiltonian cycles.

Finally, Horák and Továrek [158] considered the problem of hamiltonian cycles in complete k -partite graphs. Using this, they were able to prove the following:

Theorem 5.15 [93]. Let G be a complete k -partite graph with m vertices in each part. Let $H(G)$ be the number of hamiltonian cycles in G . Then

$$H(G) \geq \prod_{i=2}^m (k-m+1-i).$$

Note that in a very recent paper, Lovász [116] investigated the number of distinct hamiltonian cycles in a graph.

6. HIGHLY SYMMETRIC GRAPHS

In 1968, Lovász [116] conjectured that a graph containing a hamiltonian cycle and having several special orders and classes of automorphisms such graphs contain a hamiltonian cycle.

Sloane [148] asked if the existence of a pair of edge-disjoint hamiltonian cycles in G implied the existence of another such pair. Thomason [153] answered this positively.

Theorem 5.12 [153]. In a 4-regular graph of order $n \geq 3$, the number of pairs of edge-disjoint hamiltonian cycles in which two fixed edges lie in the same cycle is even.

Nincak [130] proved that if G contains k edge-disjoint hamiltonian cycles, then G contains at least $k(2k - 1)$ hamiltonian cycles. Thomason [153] showed the following:

Theorem 5.13 [153]. If a $2k$ -regular graph G of order $n \geq 3$ has a decomposition into k edge-disjoint hamiltonian cycles, then

1. each edge of G is in at least $3k - 2$ hamiltonian cycles,
2. G has at least $k(3k - 2)$ hamiltonian cycles, and
3. G has at least $(3k - 2)(3k - 5) \dots (7)(4)(1)$ hamiltonian decompositions.

Tomescu [157] considered this question for regular graphs.

Theorem 5.14 [157]. Let G be an m -regular graph of order $2m - k$ ($mk = 0 \pmod 2$).

1. If $k \geq 1$ and $m \geq 3k$, then each edge of G is contained in at least $(m - 1)(m - 2) \dots (m - k)$ hamiltonian cycles of G .
2. The graph G has at least $\frac{1}{2}(m!/(m - k - 1)!)$ hamiltonian cycles.

Finally, Horák and Továrek [93] studied the number of hamiltonian cycles in complete k -partite graphs. They obtained a recursive formula for such graphs. Using this, they were able to show the following:

Theorem 5.15 [93]. Let G be a graph of order n with $\beta_0(G) \geq m$. If $H(G)$ is the number of hamiltonian cycles in G , then $H(G) \leq \frac{1}{2}(k - m)! \prod_{i=2}^m (k - m + 1 - i)$.

Note that in a very recent paper, Cooper and Frieze [50] have investigated the number of distinct hamiltonian cycles in a random graph.

6. HIGHLY SYMMETRIC GRAPHS

In 1968, Lovász [116] conjectured that every connected vertex-transitive graph contained a hamiltonian path. This conjecture has been verified for several special orders and classes, and except for a few notable exceptions, such graphs contain a hamiltonian cycle. Babai (see [39] or [111]) proved this

conjecture for graphs with prime order $p > 2$. This follows from the work of Turner [158]. Babai [14] also showed that connected vertex-transitive graphs of order $n \geq 4$ always contain a cycle of length at least $(3n)^{1/2}$. Alspach [4] showed that every connected vertex-transitive graph of order $2p$ contained a hamiltonian cycle, unless the graph is the Petersen graph. Marusic [117] has shown that every connected vertex-transitive graph of order $p^2, p^3, 2p^2$, or $3p$ have a hamiltonian cycle; while Marusic and Parsons [118] showed graphs of order $5p$ (and $4p$) have a hamiltonian path.

Babai [13] raised the problem of constructing an infinite family of connected vertex-transitive graphs that are nonhamiltonian. To date, only a few such graphs have been found. The Petersen graph, the Coxeter graph, and the two graphs obtained from these by replacing each vertex by a triangle are the simplest such graphs. Thomassen (see [23]) conjectures that there are only finitely many such graphs.

Lipman [111] took a different approach. He considered graphs with a certain automorphism group, rather than a certain order. Let $\text{Aut } G$ denote the full automorphism group of the graph G and let Γ be a group of permutations on $V(G)$. We say Γ acts transitively if Γ has only one orbit. Using this approach Lipman was able to obtain a stronger general result.

Theorem 6.1 [111].

1. Let $\Gamma \leq \text{Aut } G$ be transitive on $V(G)$ and nilpotent. Then G has a hamiltonian path.
2. If G is a connected vertex-transitive graph and $|V(G)| = p^k, p$ a prime, then G has a hamiltonian path.

Another interesting class of graphs are the generalized Petersen graphs, $\text{GP}(n, k)$, for $n \geq 2$ and $1 \leq k < (n/2)$, with $V = \{u_0, u_1, \dots, u_{n-1}, v_0, v_1, \dots, v_{n-1}\}$ and all edges of the form $u_i u_{i+1}, u_i v_i$ and $v_i v_{i+k}$, for $0 \leq i \leq n-1$, where all subscripts are taken modulo n .

Robertson [139] proved that $\text{GP}(n, 2)$ is hamiltonian unless $n \equiv 5 \pmod{6}$. Castagna and Prins [43] conjectured that all $\text{GP}(n, k)$ were hamiltonian except for those isomorphic to $\text{GP}(n, 2)$ for $n \equiv 5 \pmod{6}$. In [6], this conjecture was verified, provided n is sufficiently large. Finally, Alspach [5] succeeded in verifying the conjecture and extending the definition of $\text{GP}(n, k)$ to the nontrivalent case $n = 2k$, he showed that $\text{GP}(n, n/2)$ is not hamiltonian if and only if $n \equiv 0 \pmod{4}$ and $n \geq 8$.

Another related class of sparse regular graphs have proven to be a little more difficult to handle. The *odd graphs*, O_k , have as their vertex set the $(k-1)$ -element subsets of a $(2k-1)$ -element set (denote these subsets as $P_{k-1}(2k-1)$). Two vertices X and Y are adjacent in O_k if $X \cap Y = \phi$. The odd graph O_3 is isomorphic to the Petersen graph.

The *Boolean graphs*, B_k , have vertex set $V = P_{k-1}(2k-1) \cup P_k(2k-1)$ and X is adjacent to Y in B_k if $X \subset Y$. Thus, B_k is the graph formed from the middle levels of the Boolean lattice of a $(2k-1)$ -element set by identi-

fying the subsets of vertices with the same size. B_k is a subset of another.

Several interesting problems remain. We say one of these graphs has a perfect matching if it can be partitioned into hamiltonian cycles. The following conjectures are proposed.

Conjecture [125]. The graph B_k is hamiltonian.

Conjecture (Foster, see [52]). The graph O_k is hamiltonian.

Conjecture [55]. The graph B_k is hamiltonian.

To date, the graphs O_4, O_5 are hamiltonian (see [125] and [111]). The Boolean graph B_3 is hamiltonian (see [101]).

As for the *odd graphs*, O_k is hamiltonian for $k \leq 5$, while O_6 is not hamiltonian in [55], while indecomposable hamiltonian.

In [55], it was noted that B_k is hamiltonian if and only if x_1 is adjacent to x_2 .

Another interesting problem is the problem of *Boolean communication*.

Problem. Let $B_i(2k-1)$ be the i -th level of the Boolean lattice of a $(2k-1)$ -element set.

$$V(B_i(2k-1)) = P_i(2k-1)$$

and X is adjacent to Y if $X \subset Y$. Are the graphs $B_i(2k-1)$ ($i \geq 1$) hamiltonian?

R. Roth (personal communication) has shown that $B_2(2k-1)$ is hamiltonian.

Another generalization also exists. They define a *uniform subset* of a $(2k-1)$ -element set as vertices and two vertices are adjacent if they correspond to k -subsets intersecting in t elements. For n, k , and t , the uniform graph is $G(n, k, t)$. The Johnson schemes are $G(n, k, k-1)$ (see [127]). The graphs $G(2k-1, k-1, 0)$ are the *odd graphs*. The following conjecture is proposed:

follows from the work
ected vertex-transitive
length at least $(3n)^{1/2}$.
ansitive graph of order
1 is the Petersen graph.
rtex-transitive graph of
ile Marasic and Parsons
hamiltonian path.
infinite family of con-
tonian. To date, only a
aph, the Coxeter graph,
ing each vertex by a tri-
e [23]) conjectures that

dered graphs with a cer-
order. Let $\text{Aut } G$ denote
t Γ be a group of permu-
as only one orbit. Using
er general result.

nilpotent. Then G has a
h and $|V(G)| = p^k, p$ a

eralized Petersen graphs,
 $V = \{u_0, u_1, \dots, u_{n-1}, v_0,$
 $u_i v_i$ and $v_i v_{i+k}$, for $0 \leq$
 n .

hamiltonian unless $n \equiv 5 \pmod 6$.
 i, k) were hamiltonian ex-
mod 6. In [6], this conjec-
ge. Finally, Alspach [5]
ending the definition of
owed that $\text{GP}(n, n/2)$ is
 $n \geq 8$.

have proven to be a little
ave as their vertex set the
t (denote these subsets as
in O_k if $X \cap Y = \phi$. The
h.
 $P_{k-1}(2k-1) \cup P_k(2k-1)$
is the graph formed from
 -1 -element set by identi-

fying the subsets as vertices with adjacency if and only if one set is a proper subset of another.

Several interesting problems have arisen on these two classes of graphs. We say one of these graphs has a hamiltonian decomposition if its edge set can be partitioned into hamiltonian cycles or hamiltonian cycles and a perfect matching.

Conjecture [125]. The graph O_k ($k \geq 4$) has a hamiltonian decomposition.

Conjecture (Erdős, see [52]). The graph B_k ($k \geq 2$) is hamiltonian.

Conjecture [55]. The graph B_k ($k \geq 2$) has a hamiltonian decomposition.

To date, the graphs O_4, O_5 , and O_6 have been shown to have a hamiltonian decomposition (see [125]), while O_7 and O_8 have been shown to be hamiltonian (see [125] and [119], respectively).

As for the Boolean graphs, B_1, B_2 , and B_3 are easily seen to have a hamiltonian decomposition, while B_4 was shown to have such a decomposition in [101]. The Boolean graphs B_5, B_6, B_7 , and B_8 were all shown to be hamiltonian in [55], while independently Dejter [52] showed B_8 and B_9 were hamiltonian.

In [55], it was noted that B_k is isomorphic to $O_k \times K_2$, where \times here represents the weak product, that is, (x_1, y_1) is adjacent to (x_2, y_2) in $G_1 \times G_2$ if and only if x_1 is adjacent to x_2 in G_1 and y_1 is adjacent to y_2 in G_2 .

Another interesting hamiltonian problem was posed by R. Roth (personal communication).

Problem. Let $B_i(2k-1)$ be the graph obtained from symmetrically opposed levels of the Boolean Lattice of an odd ordered set. That is,

$$V(B_i(2k-1)) = P_i(2k-1) \cup P_{2k-1-i}(2k-1)$$

and X is adjacent to Y if $X \subset Y$. Which generalized Boolean graphs B_i ($i \geq 1$) are hamiltonian?

R. Roth (personal communication) conjectures that each B_i ($i \geq 1$) is hamiltonian.

Another generalization along these lines is due to Chen and Lih [45]. They define a uniform subset graph $G(n, k, t)$ to have all k -subsets of an n -set as vertices and two vertices are joined by an edge if and only if the corresponding k -subsets intersect in exactly t elements. For special values of n, k , and t , the uniform subset graphs have appeared under various names. The Johnson schemes $J(n, k)$ in the theory of association schemes is $G(n, k, k-1)$ (see [127]). Kneser's graph (see [113]) is $G(2n+k, n, 0)$, while $G(2k-1, k-1, 0)$ are the odd graphs. Chen and Lih make the following conjecture:

Conjecture [45]. The graph $G(n, k, t)$ is hamiltonian for any admissible (n, k, t) except $(5, 2, 0)$ and $(5, 3, 1)$.

Heinrich and Wallis [90] proved the following:

1. The graph $G(n, k, 0)$ is hamiltonian if $n \geq k + (k2^{1/k}/(2^{1/k} - 1))$.
2. The graph $G(n, k, 0)$ is hamiltonian for
 - (a) $k = 1, n \geq 3$;
 - (b) $k = 2, n \geq 6$;
 - (c) $k = 3, n \geq 7$.

Chen and Lih [45] settle their conjecture for the cases $(n, k, k - 1)$, $(n, k, k - 2)$, and $(n, k, k - 3)$, as well as for suitably large n when k is given and t equals 0 or 1. This is not strong enough, however, to help with the odd graph conjecture.

Yet another interesting class of graphs defined from products are the hypercubes H_k , where $H_k = H_{k-1} \times K_2$ and where $H_1 = K_2$ (note that here \times denotes the usual Cartesian product). It has long been known that H_k is hamiltonian, when $k \geq 2$. However, it was conjectured that the hypercubes actually had a hamiltonian decomposition. That this is true is a consequence of a more general result of Aubert and Schneider [12].

Theorem 6.2 [12]. Let C be a cycle and let G be a graph whose edge set can be decomposed into 2 hamiltonian cycles. Then $G \times C$ (Cartesian product) can be decomposed into 3 hamiltonian cycles.

Corollary 6.3 [12].

- a. The graph $C_r \times C_s \times C_t$ is decomposable into 3 hamiltonian cycles.
- b. The graph $K_{2s+1} \times K_{2s+1} \times K_{2s+1}$ is decomposable into $3s$ hamiltonian cycles.
- c. The graph $K_{2r} \times K_{2r} \times K_{2r}$ is decomposable into $3r - 2$ hamiltonian cycles.

Let S generate the group Γ . Define the Cayley graph $\text{Cay}_s(\Gamma)$ as follows: The vertex set V corresponds to the elements of Γ and (x, xs) is an arc of $\text{Cay}_s(\Gamma)$ with initial vertex x and terminal vertex xs whenever $x \in \Gamma$ and $s \in S$. Several natural problems concerning Cayley graphs have been studied.

Problems.

1. For what generating sets S does the group Γ have a hamiltonian Cayley graph?
2. Which groups Γ have the property that for all generating sets S for Γ , $\text{Cay}_s(\Gamma)$ contains a hamiltonian path?

A great deal of work has been done in this area. Witte and Gallian [163] wrote an excellent survey article on this subject. The interested reader should begin there.

7. MISCELLANEOUS TOPIC

In this section I will consider will by no means exhaust such topics. Rather, I hope merely to and the many possible questions.

Hamiltonian properties of a in detail. In particular, Teich properties in detail.

Other graph valued functions of graphs lend themselves to higher the power (up to the d Powers of graphs were studied

Given a connected graph G

G, L^i

where $L^i(G) = L(L^{i-1}(G))$, the eventually become hamiltonian is called the hamiltonian. studying not only the hamiltonian and hamiltonian.

Many results related to the has most recently studied their relation to hamiltonian result on hamiltonian-connecte

Theorem 7.1 [167]. If G is connected.

Another special class that the following: A graph G is s $tex v$, if a hamiltonian path exists a recent survey of results on

Another strong hamiltonian through specified edges or v k -connected ($k \geq 2$), $F = \{e_1, \{e_1, \dots, e_k\}$ is connected when the edges of F . In [115], he pr Thomassen [85] proved a we graph to be $(k + 1)$ -connecte

Theorem 7.2 [85].

- (i) If L is a set of k indep incident with L are c then G has a cycle con

7. MISCELLANEOUS TOPICS

In this section I will consider several special hamiltonian problems. These will by no means exhaust such topics or even the results known on these topics. Rather, I hope merely to indicate the diversity of problems available and the many possible questions still to be asked.

Hamiltonian properties of a variety of graph products have been studied in detail. In particular, Teichert (see [149, 150, 151, 152]) has studied these properties in detail.

Other graph valued functions also can be studied. For example, powers of graphs lend themselves naturally to hamiltonian problems since the higher the power (up to the diameter), the more dense the graph becomes. Powers of graphs were studied by Paoli [134].

Given a connected graph G , if we consider the sequence of graphs

$$G, L(G), L^2(G), L^3(G), \dots$$

where $L^i(G) = L(L^{i-1}(G))$, then for $G \neq P_k$, the graphs in this sequence eventually become hamiltonian. The minimum i such that $L^i(G)$ is hamiltonian is called the *hamiltonian index* of G . Clark and Wormald [49] suggest studying not only the hamiltonian index, but similar concepts for edge-hamiltonian and hamiltonian-connected line graphs.

Many results related to the hamiltonian index have appeared. Lai [106] has most recently studied this topic. He also considered contractions and their relation to hamiltonian line graphs in [105]. Zhan [167] provided a result on hamiltonian-connected line graphs.

Theorem 7.1 [167]. If G is 4-edge connected, then $L(G)$ is hamiltonian-connected.

Another special class that has received considerable attention recently is the following: A graph G is said to be *hamiltonian-connected from a vertex* v , if a hamiltonian path exists from v to every other vertex $w \neq v$. In [51], a recent survey of results on such graphs is given.

Another strong hamiltonian property involves the existence of cycles through specified edges or vertices. Lovász [114] conjectured that if G is k -connected ($k \geq 2$), $F = \{e_1, \dots, e_k\}$ are independent edges of G and $G - \{e_1, \dots, e_k\}$ is connected when k is odd, then G contains a cycle using all the edges of F . In [115], he proved this conjecture for $k = 3$. Häggkvist and Thomassen [85] proved a weakened form of this conjecture requiring the graph to be $(k + 1)$ -connected.

Theorem 7.2 [85].

- (i) If L is a set of k independent edges in G such that any two vertices incident with L are connected by $k + 1$ internally disjoint paths, then G has a cycle containing all edges of L .

- (ii) If G is a $(\beta_0 + k)$ -connected graph, then any set of k independent edges of G is contained in a cycle.

Conjecture [85]. If G is a $\beta_0(G)$ -connected graph and L is a set of independent edges such that $G - L$ is connected, then G has a cycle containing all edges of L .

Thomassen [155] also showed that there exists a function $f(k)$ such that every strongly $f(k)$ -connected tournament has a hamiltonian cycle through any k specified edges.

Häggkvist [84] also studied a related problem. We say G is *F-hamiltonian* (*F-semihamiltonian*) if

- (i) F is a set of independent paths, and
 (ii) F is contained in a hamiltonian cycle (path).

Theorem 7.3 [84]. Let F be a 1-factor of G .

- (i) If G satisfies $\sigma_2 \geq n + 1$, then G is F -hamiltonian.
 (ii) If G satisfies $\sigma_2 \geq n - 1$, then G is F -semihamiltonian.

Häggkvist [84] also studied the degree sum of pairs of edges (another generalized degree approach) in relation to F -hamiltonian graphs. The reader interested in this should also see Woodall [164]. Cycles and paths through specified vertices have also been studied. Here I shall mention only the following: Bondy and Lovász [38] proved that a $(k + 1)$ -connected nonbipartite graph contains an odd cycle through any k specified vertices. Locke [112] showed that in an $(r + 2)$ -connected graph G with $\delta(G) \geq d$ and $|V(G)| \geq 2d - r$, any path Q of length r and any vertex y not on Q are contained in a cycle of length at least $2d - r$. In [56] the following were shown:

Theorem 7.4 [56]. Let G be a k -connected ($k \geq 2$) graph with $\delta(G) \geq d$ and order at least $2d$. Let X be a set of k vertices of G . Then G has a cycle C of length at least $2d$ such that every vertex of X is on C .

Theorem 7.5 [56]. Let G be a k -connected graph ($k \geq 3$) with $\delta(G) \geq d$ and order at least $2d - 1$. Let x and z be vertices of G and Y be a subset of $k - 1$ vertices of G . Then G has an $x - z$ path P of length at least $2d - 2$ such that every vertex of Y is on P .

Tutte [160] showed that all 4-connected planar graphs are hamiltonian. Tutte [159] also showed that some 3-connected planar graphs are nonhamiltonian. Horton (see [39]) and Ellingham and Horton [57] have constructed nonhamiltonian bipartite cubic 3-connected graphs. However, a long-standing conjecture remains.

Barnette's Conjecture [see 39, p] planar graph is hamiltonian.

In [92], some results lending credence. In particular, all such graphs are hamiltonian. They also provide

ACKNOWLEDGMENT

I would like to thank the referees for their suggestions, and to acknowledge support from grant 88-K-0070.

References

- [1] T. Agano, T. Nishizeki, Length of a hamiltonian cycle in a graph. *Graph Theory* 4 (1980) 315-336.
- [2] A. Ainouche and N. Chahboun, Hamiltonian circuits in a graph. *Math Soc. (2)* 32 (1985) 3-6.
- [3] A. Ainouche and N. Chahboun, A graph and the existence of a cycle. *Math* 17 (1987) 213-221.
- [4] B. Alspach, Hamiltonian cycles in a graph. *2p. Utilitas Math.* (1979).
- [5] B. Alspach, The classification of graphs. *J. Combinat. Theory* 17 (1984) 1-11.
- [6] B. Alspach, P. J. Robinson, Hamiltonian cycles in generalized graphs. *Math* 17 (1981) 225-231.
- [7] D. Amar, I. Fournier, Hamiltonian cycles in a graph with a given number of vertices. *International Conference on Graph Theory and Combinatorics, Israel* (1980).
- [8] D. Angluin and L. G. Valiant, Hamiltonian circuits and paths in a graph. *Math* 15 (1978) 155-193.
- [9] T. Asano, S. Kikuchi, Hamiltonian cycles in a graph. *Appl. Math* 7 (1984) 1-11.
- [10] A. S. Asratyan and N. I. Gerasimov, Hamiltonian graphs. *Mat. Zametki* 27 (1980) 1-11.

Barnette's Conjecture [see 39, p. 248]. Every cubic 3-connected bipartite planar graph is hamiltonian.

In [92], some results lending support to Barnette's conjecture are discussed. In particular, all such graphs of order at most 66 are shown to be hamiltonian. They also provide further references to related work.

ACKNOWLEDGMENT

I would like to thank the referees for their valuable and thoughtful suggestions, and to acknowledge support by ONR contract number N000014-88-K-0070.

References

- [1] T. Agano, T. Nishizeki, and T. Watanabe, An upper bound on the length of a hamiltonian walk of a maximal planar graph. *J. Graph Theory* 4 (1980) 315-336.
- [2] A. Ainouche and N. Christofides, Conditions for the existence of Hamiltonian circuits in graphs based on vertex degrees. *J. London Math Soc. (2)* 32 (1985) 385-391.
- [3] A. Ainouche and N. Christofides, Semi-independence number of a graph and the existence of Hamiltonian circuits. *Discrete Applied Math* 17 (1987) 213-221.
- [4] B. Alspach, Hamiltonian cycles in vertex-transitive graphs of order $2p$. *Utilitas Math.* (1979), 131-139.
- [5] B. Alspach, The classification of Hamiltonian generalized Petersen graphs. *J. Combinat. Theory Ser. B.* 34 (1983) 293-312.
- [6] B. Alspach, P. J. Robinson, and M. Rosenfeld, A result on Hamiltonian cycles in generalized Petersen graphs. *J. Combinat. Theory B*, 31 (1981) 225-231.
- [7] D. Amar, I. Fournier, A. Germa, and R. Häggkvist, Covering the vertices of a simple graph with given connectivity and stability number. International Conference on Convexity and Graph Theory, Israel (1980).
- [8] D. Angluin and L.G. Valiant, Fast probabilistic algorithms for Hamiltonian circuits and matchings. *J. Comput. System Sci.* 18 (1979) 155-193.
- [9] T. Asano, S. Kikuchi, and N. Saito, A linear algorithm for finding Hamiltonian cycles in 4-connected maximal-planar graphs. *Discrete Appl. Math* 7 (1984) 1-15.
- [10] A. S. Asratyan and N. K. Khachatryan, Two theorems on hamiltonian graphs. *Mat. Zametki* 35 (1984) 55-61.

- [11] A. S. Asratyan and N. K. Khachatryan, On a stability of some properties of a graph. Preprint.
- [12] J. Aubert and B. Schneider, Decomposition de la somme cartesienne d'un cycle et de l'union de deux cycles hamiltoniens en cycles hamiltoniens. *Discrete Math.* **38** (1982) 7-16.
- [13] L. Babai, Problem 17, unsolved problems. Summer Research Workshop in Algebraic Combinatorics, Simon Fraser University, July (1979).
- [14] L. Babai, Long cycles in vertex-transitive graphs. *J. Graph Theory* **3** (1979) 301-304.
- [15] D. Bauer, H. J. Broersma, and H. J. Veldman, On generalizing a theorem of Jung. Preprint.
- [16] D. Bauer, H. J. Broersma, H. J. Veldman, and L. Rao, A generalization of a result of Häggkvist and Nicoghossian. *J. Combinat. Theory B* **47** (1989) 237-243.
- [17] D. Bauer, S. L. Hamilton, and E. Schmeichel, Recognizing tough graphs is NP-hard. *Discrete Math.*, to appear.
- [18] D. Bauer, A. Morgana, E. Schmeichel, and H. J. Veldman, Long cycles in graphs with large degree sums. *Discrete Math.* **79** (1990) 59-70.
- [19] D. Bauer, E. Schmeichel, and H. J. Veldman, Some recent results on long cycles in tough graphs. *Proceedings of the Sixth International Conference on the Theory and Applications of Graphs*, to appear.
- [20] A. Benhocine, L. Clark, N. Köhler, and H. J. Veldman, On circuits and pancyclic line graphs. *J. Graph Theory* **10** (1986) 411-425.
- [21] A. Benhocine and J. Fouquet, The Chvátal-Erdős condition and pancyclic line-graphs. *Discrete Math.* **66** (1987) 21-26.
- [22] A. Benhocine and A. P. Wojda, The Geng-Hua Fan conditions for pancyclic or Hamilton-connected graphs. *J. Combinat. Theory B* **42** (1987) 167-180.
- [23] J. C. Bermond, Hamiltonian graphs. *Selected Topics in Graph Theory*. Academic Press, London (1978).
- [24] J. C. Bermond and C. Thomassen, Cycles in digraphs—A survey. *J. Graph Theory* **5** (1981) 1-43.
- [25] A. Bigulke and H. S. Jung, Über Hamiltonsche kreise und unabhängige ecken in graphen. *Monatsh. Math.* **88** (1979) 195-210.
- [26] B. Bollobás, Almost all regular graphs are Hamiltonian. *Eur. J. Combinat.* **4** (1983) 99-106.
- [27] B. Bollobás, Long paths in sparse random graphs. *Combinatorica* **2** (1982) 223-228.
- [28] B. Bollobás, *Random Graphs*. Academic Press, London (1985).
- [29] B. Bollobás, T. I. Fenner, and A. M. Frieze, An algorithm for finding Hamilton cycles in random graphs. *Proceedings of the ACM Symposium on the Theory of Computing* **17** (1985) 430-439.
- [30] B. Bollobás and R. Häggkvist, Powers of Hamiltonian cycles in tournaments. *J. Combinat. Theory B*, to appear.
- [31] J. A. Bondy, Hamilton cycles. *J. Graph Theory* **21** (1978) 3-28.
- [32] J. A. Bondy, Longest path search Report CORR 80-1, Optimization, Faculty of Mathematics, University of Toronto, Ontario, Canada (1980).
- [33] J. A. Bondy and V. Chvátal, Tough graphs. *Math. Mag.* **15** (1976) 111-136.
- [34] J. A. Bondy and G. Fan, Tough graphs. *Discrete Math.* **67** (1987) 1-10.
- [35] J. A. Bondy and R. Häggkvist, Tough graphs. *J. Graph Theory* **4** (1976) 41-44.
- [36] J. A. Bondy and A. W. H. Chan, Tough graphs. *J. Combinat. Theory B* **20** (1976) 41-44.
- [37] J. A. Bondy and K. Koulik, Tough graphs. *J. Combinat. Theory B* **42** (1987) 167-180.
- [38] J. A. Bondy and L. Lovász, Tough graphs. *Combinatorica* **1** (1981) 1-10.
- [39] J. A. Bondy and U. S. F. Leighton, Tough graphs. *J. Combinat. Theory B* **42** (1987) 167-180.
- [40] J. Bosak, Hamiltonian line graphs. *International Symposium, Rostov* (1967) 35-46.
- [41] H. J. Broersma and H. J. Veldman, Tough graphs. *J. Graph Theory* **10** (1986) 411-425.
- [42] R. A. Brualdi and R. F. Foy, Tough graphs. *J. Combinat. Theory B* **42** (1987) 167-180.
- [43] F. Castagna and G. P. Tait coloring. *Pacific J. Math.* **42** (1973) 167-180.
- [44] G. Chartrand and L. I. White, Tough graphs. *J. Graph Theory* **10** (1986) 411-425.
- [45] B. L. Chen and K. W. G. J. Combinat. Theory B **42** (1987) 167-180.
- [46] V. Chvátal, Tough graphs. *J. Graph Theory* **7** (1973) 215-228.
- [47] V. Chvátal and P. Erdős, Tough graphs. *Math. Mag.* **2** (1972) 111-113.
- [48] B. N. Clark, C. J. Colbourn, Tough graphs. *Congr. Numer. Math.* **15** (1985) 430-439.
- [49] L. H. Clark and N. C. Wormald, Tough graphs. *Ars Combinat.* **15** (1985) 430-439.
- [50] C. Cooper and A. M. Frieze, Tough graphs. *J. Graph Theory* **10** (1986) 411-425.
- [51] A. M. Dean, C. J. Knierim, Tough graphs. *J. Graph Theory* **10** (1986) 411-425.

- [31] J. A. Bondy, Hamilton cycles in graphs and digraphs. *Congr. Numer.* **21** (1978) 3–28.
- [32] J. A. Bondy, Longest paths and cycles in graphs of high degree. Research Report CORR 80-16, Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, Canada (1980).
- [33] J. A. Bondy and V. Chvátal, A method in graph theory. *Discrete Math.* **15** (1976) 111–136.
- [34] J. A. Bondy and G. Fan, A sufficient condition for dominating cycles. *Discrete Math.* **67** (1987) 205–208.
- [35] J. A. Bondy and R. Häggkvist, Edge-disjoint Hamilton cycles in 4-regular planar graphs. *Aequationes Math.* **22** (1981) 41–45.
- [36] J. A. Bondy and A. W. Ingleton, Pancyclic graphs II. *J. Combinat. Theory B* **20** (1976) 41–46.
- [37] J. A. Bondy and K. Kouider, Hamilton cycles in regular 2-connected graphs. *J. Combinat. Theory B* **44** (1988) 177–186.
- [38] J. A. Bondy and L. Lovász, Cycles through specified vertices of a graph. *Combinatorica* **1** (1981) 117–140.
- [39] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*. Macmillan & Co., London (1976).
- [40] J. Bosak, Hamiltonian lines in cubic graphs. *Theory of Graphs (International Symposium, Rome, 1966)*, Gordon and Breach, New York (1967) 35–46.
- [41] H. J. Broersma and H. J. Veldman, Restrictions on induced subgraphs ensuring hamiltonicity or pancyclicity of $K_{1,3}$ -free graphs. Preprint.
- [42] R. A. Brualdi and R. F. Shanny, Hamiltonian line graphs. *J. Graph Theory* **5** (1981) 307–314.
- [43] F. Castagna and G. Prins, Every generalized Petersen graph has a Tait coloring. *Pacific J. Math.* **40** (1972) 53–58.
- [44] G. Chartrand and L. Lesniak, *Graphs and Digraphs*. Wadsworth & Brooks/Cole, Belmont CA (1986).
- [45] B. L. Chen and K. W. Lih, Hamiltonian uniform subset graphs. *J. Combinat. Theory B* **42** (1987) 257–263.
- [46] V. Chvátal, Tough graphs and hamiltonian circuits. *Discrete Math.* **5** (1973) 215–228.
- [47] V. Chvátal and P. Erdős, A note on hamiltonian circuits. *Discrete Math.* **2** (1972) 111–113.
- [48] B. N. Clark, C. J. Colburn, and P. Erdős, A conjecture on dominating cycles. *Congr. Numer.* **47** (1985) 189–197.
- [49] L. H. Clark and N. C. Wormald, Hamiltonian-like indices of graphs. *Ars Combinat.* **15** (1983) 131–148.
- [50] C. Cooper and A. M. Frieze, On the number of Hamilton cycles in a random graph. *J. Graph Theory* **13** (1989) 719–735.
- [51] A. M. Dean, C. J. Knickerbocker, P. F. Lock, and M. Sheard, A survey of graphs hamiltonian-connected from a vertex. Preprint.

- [52] I. J. Dejter, Hamilton cycles in bipartite reflective Kneser graphs. Preprint.
- [53] G. A. Dirac, Some theorems on abstract graphs. *Proc. London Math. Soc.* 2 (1952) 69–81.
- [54] D. A. Duffus, R. J. Gould, and M. S. Jacobson, Forbidden subgraphs and the Hamiltonian theme. *Theory and Applications of Graphs* (Kalamazoo, Michigan, 1980), Wiley, New York (1981) 297–316.
- [55] D. A. Duffus, P. Hanlon, and R. Roth, Matchings and Hamiltonian cycles in some families of symmetric graphs. *J. Combinat. Theory B*, to appear.
- [56] Y. Egawa, R. Gals, and S. C. Locke, Cycles and paths through specified vertices in k -connected graphs. Preprint.
- [57] M. N. Ellingham and J. D. Horton, Non-Hamiltonian 3-connected cubic bipartite graphs. *J. Combinat. Theory Ser. B* 34 (1983) 350–353.
- [58] H. Enomoto, B. Jackson, P. Katerinis, and A. Saito, Toughness and the existence of k -factors. *J. Graph Theory* 9 (1985) 87–95.
- [59] G. H. Fan, New sufficient conditions for cycles in graphs. *J. Combinat. Theory Ser. B* 37 (1984) 221–227.
- [60] R. J. Faudree, R. J. Gould, M. S. Jacobson, and L. Lesniak, Neighborhood closures for graphs. *Colloq. Math. Societ. Janos Bolyai* 52 (1987) 227–237.
- [61] R. J. Faudree, R. J. Gould, M. S. Jacobson, and L. Lesniak, On a generalization of Dirac's theorem. *Discrete Math.*, to appear.
- [62] R. J. Faudree, R. J. Gould, M. S. Jacobson, and R. H. Schelp, Neighborhood unions and Hamiltonian properties in graphs. *J. Combinat. Theory B* 46 (1989) 1–20.
- [63] R. J. Faudree, R. J. Gould, and T. Lindquister, Hamiltonian properties and adjacency conditions in $K_{1,3}$ -free graphs. *Proceedings of the 6th International Conference on Theory and Applications of Graphs, Kalamazoo, 1988*, to appear.
- [64] R. J. Faudree, R. J. Gould, and R. H. Schelp, Neighborhood conditions and edge disjoint Hamiltonian cycles. *Congr. Numer.* 59 (1987) 55–68.
- [65] R. J. Faudree, C. C. Rousseau, and R. H. Schelp, Edge disjoint Hamiltonian cycles. *Graph Theory with Applications to Algorithms and Computer Science* (Kalamazoo, Michigan, 1984), Wiley—Interscience, New York (1985) 231–249.
- [66] T. I. Fenner and A. M. Frieze, On the existence of Hamiltonian cycles in a class of random graphs. *Discrete Math.* 45 (1983) 301–305.
- [67] T. I. Fenner and A. M. Frieze, Hamilton cycles in random graphs. *J. Combinat. Theory B* 37 (1984) 103–112.
- [68] E. Flandrin, H. A. Jung, and H. Li, Hamiltonism, degree sums and neighborhood intersections. Preprint.
- [69] E. Flandrin and H. Li, Hamiltonism and claws. Report de Recherche no. 398, Universite de Paris-Sud, Centre d'Orsay (1988).
- [70] E. Flandrin and H. Li, Hamiltonian cycles in bipartite graphs. Report de Recherche no. 40 (1988).
- [71] H. Fleischner, The square of a Hamiltonian graph. *J. Combinat. Theory*
- [72] H. Fleischner, In the square of a Hamiltonian-connected graph, Hamiltonian-connected concepts. *Monatsh. Math. Phys.*
- [73] P. Fraisse, Hamiltonian cycles and paths in bipartite graphs. Preprint.
- [74] P. Fraisse, A new sufficient condition for Hamiltonian cycles. (1986) 405–409.
- [75] A. M. Frieze, Finding Hamiltonian cycles in random bipartite graphs. *J. Combinat. Theory B* 44 (1979) 1–10.
- [76] A. M. Frieze, Limit distribution of the number of Hamiltonian cycles in random bipartite graphs. *J. Combinat. Theory B* 44 (1979) 11–18.
- [77] S. Goodman and S. Hedetniemi, Hamiltonian cycles in bipartite graphs. *J. Combinat. Theory B* 44 (1979) 19–26.
- [78] R. J. Gould, Traceability in bipartite graphs. *J. Combinat. Theory B* 44 (1979) 27–34.
- [79] R. J. Gould and M. S. Jacobson, Hamiltonian properties in graphs. *J. Combinat. Theory B* 44 (1979) 35–42.
- [80] R. J. Gould and M. S. Jacobson, Hamiltonian properties in the square of a bipartite graph. (1984) 147–154.
- [81] R. J. Gould and M. S. Jacobson, A generalization of Ore's theorem. *Conference in Graph Theory and Applications, San Francisco, 1984*.
- [82] E. J. Grinberg, Plane Hamiltonian circuits. *Lat. Am. Math. J.*
- [83] Y. Gurevich and S. Shelah, Hamiltonian path problem. *SIAM J. Comput.*
- [84] R. Häggkvist, On F -Hamiltonian graphs. *Academic Press, Topics*.
- [85] R. Häggkvist and C. Thomason, Hamiltonian cycles in bipartite graphs. *Discrete Math.* 41 (1982) 1–10.
- [86] R. Häggkvist and G. G. Lovász, Hamiltonian cycles in bipartite graphs. *J. Combinat. Theory B* 46 (1989) 1–20.
- [87] S. Hakimi and E. Schmeichel, Hamiltonian graphs. *J. Combinat. Theory B* 46 (1989) 21–28.
- [88] S. Hakimi, E. Schmeichel, Hamiltonian cycles in a bipartite graph. (1979) 365–370.
- [89] F. Harary and C. St. J. A. Bondy, Hamiltonian graphs and line graphs. *J. Combinat. Theory B* 46 (1989) 29–36.

flexive Kneser graphs.

chs. *Proc. London Math.*

on, Forbidden subgraphs
Applications of Graphs
York (1981) 297-316.

chings and Hamiltonian
J. Combinat. Theory B,

and paths through speci-
it.

Hamiltonian 3-connected
Ser. B **34** (1983) 350-353.

A. Saito, Toughness and
9 (1985) 87-95.

cles in graphs. *J. Combi-*

and L. Lesniak, Neigh-
Societ. Janos Bolyai **52**

and L. Lesniak, On a
Math., to appear.

and R. H. Schelp, Neigh-
s in graphs. *J. Combinat.*

ster, Hamiltonian proper-
graphs. *Proceedings of the*
d Applications of Graphs,

elp, Neighborhood condi-
Congr. Numer. **59** (1987)

chelp, Edge disjoint hamil-
ns to *Algorithms and Com-*
84, Wiley-Interscience,

existence of Hamiltonian
e Math. **45** (1983) 301-305.
cycles in random graphs.

iltonism, degree sums and

nd claws. Report de Re-
Centre d'Orsay (1988).

- [70] E. Flandrin and H. Li, Hamiltonism and neighborhood intersections. Report de Recherche no. 406, Universite de Paris-Sud, Centre d'Orsay (1988).
- [71] H. Fleischner, The square of every two-connected graph is hamiltonian. *J. Combinat. Theory B* **16** (1974) 29-34.
- [72] H. Fleischner, In the square of graphs, Hamiltonicity and pancyclicity, Hamiltonian-connectedness and panconnectedness are equivalent concepts. *Monatsh. Math.* **82** (1976) 125-149.
- [73] P. Fraisse, D_n -cycles and their applications for Hamiltonian graphs. Preprint.
- [74] P. Fraisse, A new sufficient condition for graphs. *J. Graph Theory* **10** (1986) 405-409.
- [75] A. M. Frieze, Finding Hamilton cycles in sparse random graphs. *J. Combinat. Theory B* **44** (1988) 230-250.
- [76] A. M. Frieze, Limit distributions for the existence of Hamilton cycles in random bipartite graphs. *Eur. J. Combinat.* **6** (1985) 327-334.
- [77] S. Goodman and S. Hedetniemi, Sufficient conditions for a graph to be hamiltonian. *J. Combinat. Theory B* **16** (1974) 175-180.
- [78] R. J. Gould, Traceability in graphs. Doctoral thesis, Western Michigan University (1979).
- [79] R. J. Gould and M. S. Jacobson, Forbidden subgraphs and Hamiltonian properties in graphs. *Discrete Math.* **42** (1982) 189-196.
- [80] R. J. Gould and M. S. Jacobson, Forbidden subgraphs and Hamiltonian properties in the square of a connected graph. *J. Graph Theory* **8** (1984) 147-154.
- [81] R. Gould and M. S. Jacobson, Neighborhood intersections and a generalization of Ore's theorem. *Proceedings of the 2nd International Conference in Graph Theory, Combinatorics, Algorithms, and Applications, San Francisco, July, 1989*, to appear.
- [82] E. J. Grinberg, Plane homogeneous graphs of degree three without hamiltonian circuits. *Latvian Math. Yearbook* **4** (1968) 51-58.
- [83] Y. Gurevich and S. Shelah, Expected computation time for Hamiltonian path problem. *SIAM J. Comput.* **16** (1987) 486-502.
- [84] R. Häggkvist, On F -Hamiltonian graphs. *Graph Theory and Related Topics*, Academic Press, New York (1979) 219-231.
- [85] R. Häggkvist and C. Thomassen, Circuits through specified edges. *Discrete Math.* **41** (1982) 29-34.
- [86] R. Häggkvist and G. G. Nicoghossian, A remark on Hamiltonian cycles. *J. Combinat. Theory B* **30** (1981) 118-120.
- [87] S. Hakimi and E. Schmeichel, A cycle structure theorem for hamiltonian graphs. *J. Combinat. Theory B* **44** (1988).
- [88] S. Hakimi, E. Schmeichel, and C. Thomassen, On the number of Hamiltonian cycles in a maximal planar graph. *J. Graph Theory* **3** (1979) 365-370.
- [89] F. Harary and C. St. J. A. Nash-Williams, On eulerian and Hamiltonian graphs and line graphs. *Can. Math. Bull.* **8** (1965) 701-710.

- [90] K. Heinrich and W.D. Wallis, Hamiltonian cycles in certain graphs. *J. Austral. Math. Soc. A* **26** (1978) 89–98.
- [91] G.R.T. Hendry and W. Vogler, The square of a connected $S(K_{1,3})$ -free graph is vertex pancyclic. *J. Graph Theory* **9** (1985) 535–537.
- [92] D.A. Holton, B. Manvel, and B.D. McKay, Hamiltonian cycles in cubic 3-connected bipartite planar graphs. *J. Combinat. Theory Ser. B* **38** (1985) 279–297.
- [93] P. Horak and L. Tovarek, On hamiltonian cycles of complete n -partite graphs (Russian summary), *Math. Slovaca* **29** (1979) 43–47.
- [94] B. Jackson, Edge-disjoint Hamilton cycles in regular graphs of large degree. *J. London Math. Soc. (2)* **19** (1979) 13–16.
- [95] B. Jackson, Hamiltonian cycles in regular 2-connected graphs. *J. Combinat. Theory B* **29** (1980) 27–46.
- [96] B. Jackson, A Chvátal-Erdős condition for Hamilton cycles in digraphs. *J. Combinat. Theory B* **43** (1987) 245–252.
- [97] B. Jackson and O. Ordaz, Chvátal-Erdős conditions for 2-cyclability in digraphs. *Ars Combinat.* **25** (1988) 39–49.
- [98] H.A. Jung, On maximal circuits in finite graphs. *Ann. Discrete Math.* **3** (1978) 129–144.
- [99] H.A. Jung and C. Nara, Note on 2-connected graphs with $d(u) + d(v) > n - 4$. *Arch. Math. (Basel)* **39** (1982) 383–384.
- [100] M. Karp, The probabilistic analysis of some combinatorial search algorithms. *Algorithms and Complexity: New Directions and Recent Results*. Academic Press, New York (1976) 1–19.
- [101] H. Kierstead and W.T. Trotter, Explicit matchings in the middle levels of the Boolean lattice. *Order* **5** (1988) 163–171.
- [102] J. Komlos and E. Szemerédi, Limit distribution for the existence of hamiltonian cycles in random graphs. *Discrete Math.* **43** (1983) 55–63.
- [103] A.D. Korshunov, Solution of a problem of Erdős and Rényi on Hamilton cycles in non-oriented graphs. *Soviet Mat. Dokl.* **17** (1976) 760–764.
- [104] A.D. Korshunov, A solution of a problem of P. Erdős and A. Rényi about Hamilton cycles in nonoriented graphs. *Metody Diskr. Anal. Teoriya Upr. Syst. Sb. Trudov Novosibirsk* **31** (1977) 17–56.
- [105] H.-J. Lai, Contractions and Hamiltonian line graphs. *J. Graph Theory* **12** (1988) 11–15.
- [106] H.-J. Lai, On the Hamiltonian index. *Discrete Math.* **69** (1988) 43–53.
- [107] E.L. Lawler, J.K. Lenstra, A.H.G. Rinnooy Kan, and D.B. Shmoys, *The Traveling Salesman Problem*. Wiley—Interscience Series, New York (1986).
- [108] L. Lesniak, Neighborhood unions and graphical properties. *Proceedings of the 6th International Conference on the Theory and Application of Graphs (Kalamazoo, 1988)*, to appear.
- [109] H. Li and Y. Zhu, Edge-disjoint Hamilton cycles in graphs. *Proceedings of the 1st China-USA Conference*, to appear.
- [110] T.E. Lindquester, The conditions on Hamiltonian graphs. *J. Graph Theory* **2** (1989) 335–352.
- [111] M. Lipman, Hamiltonian graphs with abelian and nilpotent automorphism groups. *J. Graph Theory* **1** (1988) 1–10.
- [112] S.C. Locke, A generalization of the Erdős-Rényi model. *J. Graph Theory* **1** (1985) 149–159.
- [113] L. Lovász, Kneser's conjecture and bipartite graphs. *J. Combinat. Theory A* **2** (1985) 1–10.
- [114] L. Lovász, Problem 5. *Combinatorica* **5** (1985) 1–10.
- [115] L. Lovász, *Combinatorial Problems and Exercises*. Amsterdam (1979), Section 1.1.
- [116] L. Lovász, *Combinatorial Problems and Exercises*. London (1976), Chapter 1.
- [117] D. Marusic, Hamiltonian cycles in graphs. *5p. Discrete Math.* **42** (1982) 1–10.
- [118] D. Marusic and T.D. Parsonson, Hamiltonian cycles in graphs of order $4p$. *Discrete Math.* **3** (1976) 1–10.
- [119] M. Mather, The rugby problem. *J. Graph Theory* **20** (1976) 62–63.
- [120] M.M. Matthews and D. Manvel, Hamiltonian cycles in graphs. *J. Graph Theory* **1** (1988) 1–10.
- [121] M.M. Matthews and D. Manvel, Hamiltonian cycles in free graphs. *J. Graph Theory* **1** (1988) 1–10.
- [122] C.J.H. McDiarmid, Graphs with no long paths. *Appl. Probab.* **13** (1981) 1–10.
- [123] C.J.H. McDiarmid, Graphs with no long paths. *Probab.* **15** (1983) 149–158.
- [124] J. Mitchem and E. Schmeidler, A survey. *Graphs and Applications*. Interscience, New York (1985).
- [125] G.H.J. Meredith and D. Manvel, Hamiltonian cycles in graphs. *J. Combinat. Theory B* **1** (1983) 1–10.
- [126] J. Molluzzo, Toughness and Hamiltonian cycles. *Ann. NY Acad. Sci.* **3** (1976) 1–10.
- [127] A. Moon, The graphs with no long paths. *J. Combinat. Theory B* **1** (1983) 1–10.
- [128] C. St. J.A. Nash-Williams, Hamiltonian graphs with vertices of degree n . Academic Press, London (1971).
- [129] L. Nebesky and E. Wilf, Hamiltonian cycles in powers of graph. *Canad. J. Math.* **3** (1951) 1–10.
- [130] J. Nincak, An estimate for the number of Hamiltonian cycles in a multigraph. *Recent Advances in Graph Theory (Second Czechoslovak Conference)* (1975) 431–438.

- [110] T. E. Lindquester, The effects of distance and neighborhood union conditions on Hamiltonian properties in graphs. *J. Graph Theory* **13** (1989) 335–352.
- [111] M. Lipman, Hamiltonian cycles and paths in vertex-transitive graphs with abelian and nilpotent groups. *Discrete Math.* **54** (1985) 15–21.
- [112] S. C. Locke, A generalization of Dirac's theorem. *Combinatorica* **5** (1985) 149–159.
- [113] L. Lovász, Kneser's conjecture, chromatic number and homotopy. *J. Combinat. Theory A* **25** (1978) 319–324.
- [114] L. Lovász, Problem 5. *Period. Math. Hungar.* **4** (1974) 82.
- [115] L. Lovász, *Combinatorial Problems and Exercises*. North Holland, Amsterdam (1979), Section 6.67.
- [116] L. Lovász, *Combinatorial Structures and Their Applications*. Gordon and Breach, London (1970) Problem 11.
- [117] D. Marusic, Hamiltonian paths in vertex-symmetric graphs of order $5p$. *Discrete Math.* **42** (1982) 227–242.
- [118] D. Marusic and T. D. Parsons, Hamiltonian paths in vertex-symmetric graphs of order $4p$. *Discrete Math.* **43** (1983) 91–96.
- [119] M. Mather, The rugby footballers of Croam. *J. Combinat. Theory B.* **20** (1976) 62–63.
- [120] M. M. Matthews and D. P. Sumner, Hamiltonian results in $K_{1,3}$ -free graphs. *J. Graph Theory* **8** (1984) 139–146.
- [121] M. M. Matthews and D. P. Sumner, Longest paths and cycles in $K_{1,3}$ -free graphs. *J. Graph Theory* **9** (1985) 269–277.
- [122] C. J. H. McDiarmid, General percolation and random graphs. *Adv. Appl. Probab.* **13** (1981) 40–60.
- [123] C. J. H. McDiarmid, General first-passage percolation. *Adv. Appl. Probab.* **15** (1983) 149–161.
- [124] J. Mitchem and E. Schmeichel, Pancyclic and bypancyclic graphs—A survey. *Graphs and Applications (Boulder, Colorado, 1982)*. Wiley—Interscience, New York (1985) 271–287.
- [125] G. H. J. Meredith and E. K. Lloyd, The footballers of Croam. *J. Combinat. Theory B.* **15** (1973) 161–166.
- [126] J. Molluzzo, Toughness, Hamiltonian-connectedness, and n -Hamiltonicity. *Ann. NY Acad. Sci.* **319** (1979) 402–404.
- [127] A. Moon, The graphs $G(n, k)$ of the Johnson schemes are unique for $n \geq 20$. *J. Combinat. Theory B* **37** (1984) 173–188.
- [128] C. St. J. A. Nash-Williams, Edge-disjoint Hamiltonian circuits in graphs with vertices of large valency. *Studies in Pure Mathematics*, Academic Press, London (1971) 157–183.
- [129] L. Nebesky and E. Wisztova, Two edge-disjoint Hamiltonian cycles of powers of graph. *Casopi Pest. Mat.* **110** (1985) 294–301, 315.
- [130] J. Nincak, An estimate of the number of Hamiltonian cycles in a multigraph. *Recent Advances in Graph Theory (Proceedings of the Second Czechoslovak Symposium, Prague, 1974)*. Academia, Prague (1975) 431–438.

- [131] D. J. Oberly and D. P. Sumner, Every connected, locally connected nontrivial graph with no induced claw is hamiltonian. *J. Graph Theory* 3 (1979) 351–356.
- [132] O. Ore, A note on hamiltonian circuits. *Am. Math. Month.* 67 (1960) 55.
- [133] P. J. Owens, On regular graphs and Hamiltonian circuits, including answers to some questions of Joseph Zaks. *J. Combinat. Theory B* 28 (1980) 263–277.
- [134] M. Paoli, Powers of connected graphs and hamiltonicity. *Discrete Math.* 52 (1984) 91–99.
- [135] L. Pósa, A theorem concerning hamilton lines. *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 7 (1962) 225–226.
- [136] L. Pósa, Hamiltonian circuits in random graphs. *Discrete Math.* 14 (1976) 359–364.
- [137] L. B. Richmond, R. W. Robinson, and N. C. Wormald, On Hamiltonian cycles in 3-connected cubic maps. *Ann. Discrete Math.* 27 (1985) 141–150.
- [138] S. Riha, A new proof of a theorem by Fleischner. *J. Combinat. Theory B*, to appear.
- [139] G. N. Robertson, Graphs with girth, valency, and connectivity constraints. Ph.D. thesis, University of Waterloo, Ontario (1968).
- [140] R. W. Robinson and N. C. Wormald, Almost all bipartite cubic graphs are hamiltonian. Preprint.
- [141] I. Schiermeyer, A polynomial algorithm for hamiltonian graphs based on transformations. *Ars Combinat.* 25B (1988) 55–77.
- [142] I. Schiermeyer, A strong closure concept for Hamiltonian graphs. Preprint.
- [143] A. I. Serdjukov, The problem of finding a Hamilton cycle (contour) in the presence of exclusions. *Upravljaemye Sistemy* 19 (1979) 57–64, 79–80.
- [144] E. Shamir, How many random edges makes a graph Hamiltonian? *Combinatorica* 3 (1983) 123–131.
- [145] J. Sheehan, Graphs with exactly one Hamiltonian circuit. *J. Graph Theory* 1 (1977) 37–43.
- [146] J. Sheehan and E. M. Wright, The number of Hamiltonian circuits in large, heavily edged graphs. *Glasgow Math. J.* 18 (1977) 35–37.
- [147] Z. Skupien, Sharp sufficient conditions for hamiltonian cycles in tough graphs. Preprint.
- [148] N. J. A. Sloane, Hamiltonian cycles in a graph of degree 4. *J. Combinat. Theory* 6 (1969) 311–312.
- [149] H. M. Teichert, Connectivity and hamiltonian properties of the disjunction of undirected graphs. *Elektron. Informatsverab. Kybernet* 19 (1983) 611–623.
- [150] H. M. Teichert, Hamiltonian properties of the lexicographic product of undirected graphs. *Elektron. Informationsverarb. Kybernet.* 19 (1983) 67–77.
- [151] H. M. Teichert, On the *tron. Informationsverarb.*
- [152] H. M. Teichert, On the *C. Informationsverarb. Kyb.*
- [153] A. Thomason, Hamiltonian graphs. *Ann. Discrete Math.*
- [154] C. Thomassen, Long cycles. *J. Combinat. Theory B* 42 (1981) 231–251.
- [155] C. Thomassen, Connected graphs. *Combinatorics, Academic Press*
- [156] G. L. Thompson and S. Cioabă, Directed hamiltonian paths. *J. Combinat. Theory B* 179–195.
- [157] I. Tomescu, Hamiltonian graphs. *Math. Press Appl.* 29 (1980) 1–10.
- [158] J. Turner, Point-symmetric graphs. *J. Combin. Theory B* 3 (1975) 1–10.
- [159] W. T. Tutte, On Hamiltonian graphs. *Canad. J. Math.* 98–101.
- [160] W. T. Tutte, ed., *Recent Advances in Graph Theory*. New York (1969).
- [161] H. J. Veldman, Existence of hamiltonian paths. *Math.* 43 (1983) 281–294.
- [162] H. J. Veldman, Existence of hamiltonian cycles. *Math.* 44 (1983) 309–316.
- [163] D. Witte and J. A. Gallian, Hamiltonian graphs. *Discrete Math.*
- [164] D. R. Woodall, Circuits in graphs. *J. Combinat. Theory B*
- [165] J. Zaks, Pairs of Hamiltonian paths. *J. Combinat. Theory B*
- [166] J. Zaks, Extending an Hamiltonian path. *J. Graph Theory* B 32 (1982) 95–104.
- [167] S. Zhan, Hamiltonian paths in graphs. *J. Graph Theory* 22 (1986) 89–95.
- [168] C. Q. Zhang, Hamiltonian paths in graphs. *J. Graph Theory* 12 (1988) 209–216.
- [169] Y. J. Zhu, Z. H. Liu, A result on Hamiltonian cycles in graphs. *Graphs (Burnaby, B.C.)* Vol. 115. North Holland
- [170] Y. J. Zhu and F. Tian, Hamiltonian paths in graphs. *Graphs (Burnaby, B.C.)* Vol. 115. North Holland

- ted, locally connected hamiltonian. *J. Graph Theory*. **67** (1960)
- ian circuits, including hamiltonicity. *Discrete Math.* **14** (1985)
- ombinat. *Theory B* **28** (1985)
- s. *Magyar Tud. Akad. Discrete Math.* **14** (1985)
- ormald, On Hamiltonian graphs based on Hamiltonian graphs. *J. Combinat. Theory B* **3** (1967) 136-145.
- and connectivity con- Ontario (1968). bipartite cubic graphs hamiltonian graphs based on Hamiltonian graphs. *J. Combinat. Theory* **19** (1979) 57-64,
- graph Hamiltonian?
- ian circuit. *J. Graph Theory* **8** (1977) 35-37.
- amiltonian circuits in hamiltonian cycles in of degree 4. *J. Combinat. Theory* **19** (1979) 57-64,
- properties of the dis- *atsverarb. Kybernet* **19** (1983) 345-356.
- lexicographic prod- *sverarb. Kybernet*. **19** (1982) 639-646.
- [153] A. Thomason, Hamiltonian cycles and uniquely edge colourable graphs. *Ann. Discrete Math.* **3** (1978) 259-268.
- [154] C. Thomassen, Long cycles in digraphs. *Proc. London Math. Soc.* (3) **42** (1981) 231-251.
- [155] C. Thomassen, Connectivity in tournaments. *Graph Theory and Combinatorics*, Academic Press, Orlando, FL (1984) 305-313.
- [156] G. L. Thompson and S. Singhal, A successful algorithm for the undirected hamiltonian path problem. *Discrete Appl. Math.* **10** (1985) 179-195.
- [157] I. Tomescu, Hamiltonian property of regular graphs. *Rev. Roumaine Math. Press Appl.* **29** (1984) 499-505.
- [158] J. Turner, Point-symmetric graphs with a prime number of points. *J. Combinat. Theory B* **3** (1967) 136-145.
- [159] W.T. Tutte, On Hamiltonian circuits. *J. London Math. Soc.* **21** (1946) 98-101.
- [160] W.T. Tutte, ed., *Recent Progress in Combinatorics*. Academic Press, New York (1969).
- [161] H. J. Veldman, Existence of dominating cycles and paths. *Discrete Math.* **43** (1983) 281-296.
- [162] H. J. Veldman, Existence of D_λ -cycles and D_λ -paths. *Discrete Math.* **44** (1983) 309-316.
- [163] D. Witte and J. A. Gallian, A survey-Hamiltonian cycles in Cayley graphs. *Discrete Math.* **51** (1984) 293-304.
- [164] D. R. Woodall, Circuits containing specified edges of a graph. *J. Combinat. Theory B* **22** (1977) 274-278.
- [165] J. Zaks, Pairs of Hamiltonian circuits in 5-connected planar graphs. *J. Combinat. Theory B* **21** (1976) 116-131.
- [166] J. Zaks, Extending an extension of Grinberg's theorem. *J. Combinat. Theory B* **32** (1982) 95-98.
- [167] S. Zhan, Hamiltonian-connectedness of line graphs. *Ars Combinat.* **22** (1986) 89-95.
- [168] C. Q. Zhang, Hamiltonian cycles in claw-free graphs. *J. Graph Theory* **12** (1988) 209-216.
- [169] Y. J. Zhu, Z. H. Liu, and Z. G. Yu, An improvement of Jackson's result on Hamilton cycles in 2-connected regular graphs. *Cycles in Graphs* (Burnaby, B.C., 1982), North Holland Mathematics Studies, Vol. 115. North Holland, Amsterdam (1985) 237-247.
- [170] Y. J. Zhu and F. Tian, A generalization of the Bondy-Chvátal theorem on the k -closure. *J. Combinat. Theory Ser. B.* **35** (1983) 247-255.